

Nullspace of Compound Magic Squares

Saleem Al-Ashhab¹

¹Departemnet of Mathematics, Al-albayt University, P.O. Box 130040
 Mafraq / Jordan
 e-mail: ahhab@aabu.edu.jo

Abstract

In this paper we consider special compound $4n \times 4n$ and $6n \times 6n$ magic squares. We determine a $2n-3$ (res. $3n-5$) dimensional subspace of the nullspace of the $4n \times 4n$ (res. $6n \times 6n$) squares. All vectors in the subspaces possess the property that the sum of all entries of each vector equals zero.

Keywords: Null space, Magic squares, Mathematical induction.

2010 AMS classification number: Primary 15A15. Secondary 16A24, 15A30.

1 Introduction

A semi magic square is a $n \times n$ matrix such that the sum of the entries in each row and columns is the same. The common value is called the magic constant. If, in addition, the sum of all entries in each left-broken diagonal and each right-broken diagonal is the magic constant, then we call the matrix a pandiagonal magic square. Rosser and Walker show that a pandiagonal 4×4 magic square with magic constant $2s$ has in general the following structure

A	B	C	ω
E	θ	ς	ρ
$s - C$	$s - \omega$	$s - A$	$s - B$
$s - \varsigma$	$s - \rho$	$s - E$	$s - \theta$

where

$$\omega = 2s - A - B - C,$$

$$\begin{aligned}\theta &= 2s - A - B - E, \\ \varsigma &= A + E - C, \\ \rho &= B + C - E.\end{aligned}$$

We note that every antipodal pair of elements add up to one half of the magic constant.

We define a certain class of 6×6 magic squares, which has a similar structure to the structure of a pandiagonal 4×4 magic square. In this class each antipodal pair will add up to one third of the magic constant. Precisely, we have

Definition 1.1: A 6×6 magic square with $3s$ as a magic constant is called panmagic if

$$a_{ij} + a_{kl} = s, \text{ for each } i, j, k, l \text{ such that } i \equiv k \pmod{3} \text{ and } j \equiv l \pmod{3}.$$

The following matrix is a possible form for this kind of squares:

M	R	W	T	L	K
Q	J	I	H	G	F
P	E	D	C	B	A
$s - T$	$s - L$	$s - K$	$s - M$	$s - R$	$s - W$
$s - H$	$s - G$	$s - F$	$s - Q$	$s - J$	$s - I$
$s - C$	$s - B$	$s - A$	$s - P$	$s - E$	$s - D$

where

$$M = J + I + H + E + D + C - L - K - \frac{3s}{2},$$

$$W = K - I + F - D + A,$$

$$P = 3s - E - D - C - B - A$$

$$Q = 3s - J - I - H - G - F,$$

$$R = L - J + G - E + B,$$

$$T = \frac{9s}{2} - L - K - H - G - F - C - B - A.$$

Note that we have the following relations:

$$M + Q + P = T + H + C,$$

$$R + J + E = L + G + B,$$

$$W + I + D = K + F + A.$$

Using Maple we can show that the 6×6 panmagic square possesses a nontrivial null space, which can be written in the following form:

$$\{z(x_1, x_2, x_3, -x_1, -x_2, -x_3) : z \in R\}$$

where

$$\begin{aligned}x_1 &= (A - D)(G - J)(B - E)(I - F), \\x_2 &= (F - I)(B + 2C + E - 3s) + 2FD - 2AI + (D - A)(G + J + 2H - 3s), \\x_3 &= (B - E)(F + I + 2H) + (A + D + 2C + 2B + 2E - 3s)(J - G).\end{aligned}$$

Note that the sum of all entries of the vectors is zero.

For example,

-51	39	26	0	9	13
54	-10	-2	-5	4	-5
-5	1	2	3	17	18
12	3	-1	63	-27	-14
17	8	17	-42	22	14
9	-5	-6	17	11	10

has as nullspace $\{z(34, 115, -132, -34, -115, 132)^t : z \in R\}$.

Definition 1.2: A 8×8 square consisting of 4 pandiagonal magic squares $A_{11}, A_{12}, A_{21}, A_{22}$ having the same magic sum in the form

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$$

is called a compound magic square if the following relation holds

$$A_{22} + A_{11} = A_{12} + A_{21}.$$

It is easy to check that the last relation guarantees that the square is a magic 8×8 square. In the same manner we can combine four panmagic squares in a magic square.

Definition 1.3: Let $B_{22}, B_{11}, B_{12}, B_{21}$ be panmagic squares having the same magic constant. Assume that $B_{22} + B_{11} = B_{12} + B_{21}$. Then the matrix

$$\begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

is called the compound 12×12 magic square.

The condition $B_{22} + B_{11} = B_{12} + B_{21}$ ensures that the compound 12×12 magic square is magic.

2 Main Results

We prove first a simple result for a compound square of 4×4 squares. We then generalize this result for an arbitrary number of squares.

Proposition 2.1: The compound 8×8 magic square processes a three dimensional subspace of its nullspace.

Proof: First we note that the vector

$$(1,1,1,1,-1,-1,-1,-1)'$$

is a nonzero vector, which belongs to the nullspace of the square, since the squares have the same magic constant.

Now, the square A_{11} (res. A_{12}) has a nonzero vector v_{11} (res. v_{12}), which belongs to the nullspace of the square, since A_{11} (res. A_{12}) is a pandiagonal magic square.

We look for four numbers $f_{11}, f_{12}, f_{21}, f_{22}$ such that the vector

$$\begin{pmatrix} f_{11}v_{11} + f_{12}v_{12} \\ f_{21}v_{11} + f_{22}v_{12} \end{pmatrix}$$

belongs to the nullspace of the square. To do this we compute the following matrix multiplication

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{pmatrix} f_{11}v_{11} + f_{12}v_{12} \\ f_{21}v_{11} + f_{22}v_{12} \end{pmatrix} = \begin{pmatrix} A_{11}(f_{11}v_{11} + f_{12}v_{12}) + A_{12}(f_{21}v_{11} + f_{22}v_{12}) \\ A_{21}(f_{11}v_{11} + f_{12}v_{12}) + A_{22}(f_{21}v_{11} + f_{22}v_{12}) \end{pmatrix}$$

According to the choice of v_{11} and v_{12} we obtain the vector $(g_1, g_2)'$ as the result of matrix multiplication, where

$$g_1 = A_{11}f_{12}v_{12} + A_{12}f_{21}v_{11},$$

$$g_2 = A_{21}v_{11}f_{11} + A_{21}v_{12}f_{12} + (A_{12}v_{11} + A_{21}v_{11})f_{21} + (A_{21}v_{12} - A_{11}v_{12})f_{22}$$

Note that we used the relation $A_{22} = A_{12} + A_{21} - A_{11}$. We can rewrite the vector $(g_1, g_2)'$ in the form

$$\begin{bmatrix} 0 & A_{11}v_{12} & A_{12}v_{11} & 0 \\ A_{21}v_{11} & A_{21}v_{12} & (A_{12} + A_{21})v_{11} & (A_{21} - A_{11})v_{12} \end{bmatrix} \begin{pmatrix} f_{11} \\ f_{12} \\ f_{21} \\ f_{22} \end{pmatrix} \dots \dots \dots \quad (1)$$

According to Al-Ashhab we can assume that the vectors in the nullspace of the pandiagonal magic square

$$v_{ij} = (v_{ij}^*, v_{ij}^{**}, -v_{ij}^*, -v_{ij}^{**})', \text{ for } i=1, j=1, 2$$

Further, we can assume that

$$A_{ij} = \begin{bmatrix} a_{ij} & b_{ij} & c_{ij} & d_{ij} \\ e_{ij} & f_{ij} & g_{ij} & h_{ij} \\ s - c_{ij} & s - d_{ij} & s - a_{ij} & s - b_{ij} \\ s - g_{ij} & s - h_{ij} & s - e_{ij} & s - f_{ij} \end{bmatrix}, \quad i, j = 1, 2$$

Hence, we can assume that

$$A_{ij}v_{ij} = \begin{pmatrix} a_{ij}v_{ij}^* + b_{ij}v_{ij}^{**} - c_{ij}v_{ij}^* - d_{ij}v_{ij}^{**} \\ e_{ij}v_{ij}^* + f_{ij}v_{ij}^{**} - g_{ij}v_{ij}^* - h_{ij}v_{ij}^{**} \\ -c_{ij}v_{ij}^* - d_{ij}v_{ij}^{**} + a_{ij}v_{ij}^* + b_{ij}v_{ij}^{**} \\ -g_{ij}v_{ij}^* - h_{ij}v_{ij}^{**} + e_{ij}v_{ij}^* + f_{ij}v_{ij}^{**} \end{pmatrix} = \begin{pmatrix} (a_{ij} - c_{ij})v_{ij}^* + (b_{ij} - d_{ij})v_{ij}^{**} \\ (e_{ij} - g_{ij})v_{ij}^* + (f_{ij} - h_{ij})v_{ij}^{**} \\ -(a_{ij} - c_{ij})v_{ij}^* - (b_{ij} - d_{ij})v_{ij}^{**} \\ -(e_{ij} - g_{ij})v_{ij}^* - (f_{ij} - h_{ij})v_{ij}^{**} \end{pmatrix}$$

Since the sum of two pandiagonal magic squares is pandiagonal magic, we deduce that four rows in the matrix in (1) are redundant. Since we have the relations

$$a_{11} + e_{11} = c_{11} + g_{11} \Rightarrow a_{11} - c_{11} = -(e_{11} - g_{11})$$

$$b_{11} + f_{11} = d_{11} + h_{11} \Rightarrow b_{11} - d_{11} = -(f_{11} - h_{11})$$

the application of elementary row operations on the matrix in (1) yields to

$$\left[\begin{array}{cccc} 0 & r_{12} & r_{21} & 0 \\ q_{11} & q_{12} & r_{21} + q_{11} & q_{12} - r_{12} \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{array} \right]$$

where

$$r_{12} = (a_{11} - c_{11})v_{12}^* + (b_{11} - d_{11})v_{12}^{**}$$

$$r_{21} = (a_{12} - c_{12})v_{11}^* + (b_{12} - d_{12})v_{11}^{**}$$

$$q_{11} = (a_{21} - c_{21})v_{11}^* + (b_{21} - d_{21})v_{11}^{**}$$

$$q_{12} = (a_{21} - c_{21})v_{12}^* + (b_{21} - d_{21})v_{12}^{**}$$

This analysis enables us to conclude the following relations from (1)

$$f_{11} = -\frac{r_{12}r_{21} + q_{11}r_{12} - q_{12}r_{21}}{q_{11}r_{12}} f_{21} + \frac{-q_{12} + r_{12}}{q_{11}} f_{22},$$

$$f_{12} = -\frac{r_{21}}{r_{12}} f_{21}.$$

If we set

$$f_{12} = 0, f_{21} = 0, f_{22} = q_{11}, f_{11} = r_{12} - q_{12},$$

which is consistent with the previous relations, we conclude that the vector

$$\begin{pmatrix} (r_{12} - q_{12})v_{11} \\ q_{11}v_{12} \end{pmatrix}$$

belongs to the nullspace of the square. We can make another choice as follows

$$f_{22} = 0, f_{21} = r_{12}q_{11}, f_{12} = -r_{21}q_{11}, f_{11} = r_{21}q_{12} - r_{12}(r_{21} + q_{11})$$

and we obtain a vector belonging to the nullspace of the square, which is

$$\begin{pmatrix} (r_{21}q_{12} - r_{12}(r_{21} + q_{11}))v_{11} - r_{21}q_{11}v_{12} \\ -r_{21}q_{11}v_{11} \end{pmatrix}$$

Now, the vectors v_{12}, v_{11} are linearly independent, since they correspond to different magic squares. Hence, the last two vectors are linearly independent. Also the vector

$$(1,1,1,1,-1,-1,-1,-1)'$$

is linearly independent with the last two vectors, since its first two entries are not the opposite of the third and fourth entry. \square

For example, the following square is a compound 8×8 magic square

0	14	-19	13	10	5	-22	15
-12	6	7	7	-20	13	12	3
23	-9	4	-10	26	-11	-6	-1
-3	-3	16	-2	-8	1	24	-9
-16	25	-17	16	-6	16	-20	18
1	-2	2	7	-7	5	7	3
21	-12	20	-21	24	-14	10	-12
2	-3	3	6	-3	1	11	-1

For this square we can construct as described the following two vectors in its nullspace

$$\begin{pmatrix} -\frac{38}{5}, \frac{722}{5}, \frac{38}{5}, -\frac{722}{5}, -170, -544, 170, 544 \end{pmatrix}'$$

$$\begin{pmatrix} -\frac{216006}{5}, -\frac{85026}{5}, \frac{216006}{5}, \frac{85026}{5}, 3774, -71706, -3774, 71706 \end{pmatrix}'$$

In fact, its nullity is 3. Thus, these two vectors together with

$$(1,1,1,1,-1,-1,-1,-1)'$$

form a basis of its nullspace.

We prove now a similar result to the previous proposition, where we replace the 4×4 square with a 6×6 one.

Proposition 2.2: The compound 12×12 magic square possess a three dimensional subspace of its nullspace.

Proof: First we note that the vector

$$(1,1,1,1,1,-1,-1,-1,-1,-1,-1,-1)'$$

is a nonzero vector, which belongs to the nullspace of the square, since the squares have the same magic constant.

We look for scalars $v_1, v_2, v_3, v_4, v_5, v_6$ such that

$$\left[\begin{array}{ccccccccc} a_{11} & b_{11} & c_{11} & d_{11} & e_{11} & f_{11} & a_{12} & b_{12} & c_{12} \\ g_{11} & h_{11} & i_{11} & j_{11} & k_{11} & l_{11} & g_{12} & h_{12} & i_{12} \\ m_{11} & n_{11} & o_{11} & p_{11} & q_{11} & r_{11} & m_{12} & n_{12} & o_{12} \\ s - d_{11}s - e_{11}s - f_{11}s - a_{11}s - b_{11}s - c_{11}s - d_{12}s - e_{12}s - f_{12}s - a_{12}s - b_{12}s - c_{12}s \\ s - j_{11}s - k_{11}s - l_{11}s - g_{11}s - h_{11}s - i_{11}s - j_{12}s - k_{12}s - l_{12}s - g_{12}s - h_{12}s - i_{12}s \\ s - p_{11}s - q_{11}s - r_{11}s - m_{11}s - n_{11}s - o_{11}s - p_{12}s - q_{12}s - r_{12}s - m_{12}s - n_{12}s - o_{12}s \\ a_{21} & b_{21} & c_{21} & d_{21} & e_{21} & f_{21} & a_{22} & b_{22} & c_{22} \\ g_{21} & h_{21} & i_{21} & j_{21} & k_{21} & l_{21} & g_{22} & h_{22} & i_{22} \\ m_{21} & n_{21} & o_{21} & p_{21} & q_{21} & r_{21} & m_{22} & n_{22} & o_{22} \\ s - d_{21}s - e_{21}s - f_{21}s - a_{21}s - b_{21}s - c_{21}s - d_{22}s - e_{22}s - f_{22}s - a_{22}s - b_{22}s - c_{22}s \\ s - j_{21}s - k_{21}s - l_{21}s - g_{21}s - h_{21}s - i_{21}s - j_{22}s - k_{22}s - l_{22}s - g_{22}s - h_{22}s - i_{22}s \\ s - p_{21}s - q_{21}s - r_{21}s - m_{21}s - n_{21}s - o_{21}s - p_{22}s - q_{22}s - r_{22}s - m_{22}s - n_{22}s - o_{22}s \end{array} \right] \begin{bmatrix} v_1 \\ v_2 \\ v_3 \\ v_4 \\ v_5 \\ v_6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ -v_1 \\ -v_2 \\ -v_3 \\ v_4 \\ v_5 \\ v_6 \\ -v_4 \\ -v_5 \\ -v_6 \\ 0 \end{bmatrix}$$

We transform this equation into the linear system

$$\begin{aligned} a_{11}v_1 + b_{11}v_2 + c_{11}v_3 - d_{11}v_1 - e_{11}v_2 - f_{11}v_3 + a_{12}v_4 + b_{12}v_5 + c_{12}v_6 - d_{12}v_4 - e_{12}v_5 - f_{12}v_6 &= 0 \\ g_{11}v_1 + h_{11}v_2 + i_{11}v_3 - j_{11}v_1 - k_{11}v_2 - l_{11}v_3 + g_{12}v_4 + h_{12}v_5 + i_{12}v_6 - j_{12}v_4 - k_{12}v_5 - l_{12}v_6 &= 0 \\ m_{11}v_1 + n_{11}v_2 + o_{11}v_3 - p_{11}v_1 - q_{11}v_2 - r_{11}v_3 + m_{12}v_4 + n_{12}v_5 + o_{12}v_6 - p_{12}v_4 - q_{12}v_5 - r_{12}v_6 &= 0 \\ -d_{11}v_1 - e_{11}v_2 - f_{11}v_3 + a_{11}v_1 + b_{11}v_2 + c_{11}v_3 - d_{12}v_4 - e_{12}v_5 - f_{12}v_6 + a_{12}v_4 + b_{12}v_5 + c_{12}v_6 &= 0 \\ -j_{11}v_1 - k_{11}v_2 - l_{11}v_3 + g_{11}v_1 + h_{11}v_2 + i_{11}v_3 - j_{12}v_4 - k_{12}v_5 - l_{12}v_6 + g_{12}v_4 + h_{12}v_5 + i_{12}v_6 &= 0 \\ -p_{11}v_1 - q_{11}v_2 - r_{11}v_3 + m_{11}v_1 + n_{11}v_2 + o_{11}v_3 - p_{12}v_4 - q_{12}v_5 - r_{12}v_6 + m_{12}v_4 + n_{12}v_5 + o_{12}v_6 &= 0 \\ a_{21}v_1 + b_{21}v_2 + c_{21}v_3 - d_{21}v_1 - e_{21}v_2 - f_{21}v_3 + a_{22}v_4 + b_{22}v_5 + c_{22}v_6 - d_{22}v_4 - e_{22}v_5 - f_{22}v_6 &= 0 \\ g_{21}v_1 + h_{21}v_2 + i_{21}v_3 - j_{21}v_1 - k_{21}v_2 - l_{21}v_3 + g_{22}v_4 + h_{22}v_5 + i_{22}v_6 - j_{22}v_4 - k_{22}v_5 - l_{22}v_6 &= 0 \\ m_{21}v_1 + n_{21}v_2 + o_{21}v_3 - p_{21}v_1 - q_{21}v_2 - r_{21}v_3 + m_{22}v_4 + n_{22}v_5 + o_{22}v_6 - p_{22}v_4 - q_{22}v_5 - r_{22}v_6 &= 0 \\ -d_{21}v_1 - e_{21}v_2 - f_{21}v_3 + a_{21}v_1 + b_{21}v_2 + c_{21}v_3 - d_{22}v_4 - e_{22}v_5 - f_{22}v_6 + a_{22}v_4 + b_{22}v_5 + c_{22}v_6 &= 0 \\ -j_{21}v_1 - k_{21}v_2 - l_{21}v_3 + g_{21}v_1 + h_{21}v_2 + i_{21}v_3 - j_{22}v_4 - k_{22}v_5 - l_{22}v_6 + g_{22}v_4 + h_{22}v_5 + i_{22}v_6 &= 0 \\ -p_{21}v_1 - q_{21}v_2 - r_{21}v_3 + m_{21}v_1 + n_{21}v_2 + o_{21}v_3 - p_{22}v_4 - q_{22}v_5 - r_{22}v_6 + m_{22}v_4 + n_{22}v_5 + o_{22}v_6 &= 0 \end{aligned}$$

We eliminate the redundant equations. The system becomes

$$\begin{aligned} (a_{11} - d_{11})v_1 + (b_{11} - e_{11})v_2 + (c_{11} - f_{11})v_3 + (a_{12} - d_{12})v_4 + (b_{12} - e_{12})v_5 + (c_{12} - f_{12})v_6 &= 0 \\ (g_{11} - j_{11})v_1 + (h_{11} - k_{11})v_2 + (i_{11} - l_{11})v_3 + (g_{12} - j_{12})v_4 + (h_{12} - k_{12})v_5 + (i_{12} - l_{12})v_6 &= 0 \\ (m_{11} - p_{11})v_1 + (n_{11} - q_{11})v_2 + (o_{11} - r_{11})v_3 + (m_{12} - p_{12})v_4 + (n_{12} - q_{12})v_5 + (o_{12} - r_{12})v_6 &= 0 \\ (a_{21} - d_{21})v_1 + (b_{21} - e_{21})v_2 + (c_{21} - f_{21})v_3 + (a_{22} - d_{22})v_4 + (b_{22} - e_{22})v_5 + (c_{22} - f_{22})v_6 &= 0 \\ (g_{21} - j_{21})v_1 + (h_{21} - k_{21})v_2 + (i_{21} - l_{21})v_3 + (g_{22} - j_{22})v_4 + (h_{22} - k_{22})v_5 + (i_{22} - l_{22})v_6 &= 0 \\ (m_{21} - p_{21})v_1 + (n_{21} - q_{21})v_2 + (o_{21} - r_{21})v_3 + (m_{22} - p_{22})v_4 + (n_{22} - q_{22})v_5 + (o_{22} - r_{22})v_6 &= 0 \end{aligned}$$

From the definition of the panmagic square we know that

$$a_{ij} + g_{ij} + m_{ij} = d_{ij} + j_{ij} + p_{ij} \Rightarrow (a_{ij} - d_{ij}) + (g_{ij} - j_{ij}) = -(m_{ij} - p_{ij}) \dots\dots\dots(1)$$

$$b_{ij} + h_{ij} + n_{ij} = e_{ij} + k_{ij} + q_{ij} \Rightarrow (b_{ij} - e_{ij}) + (h_{ij} - k_{ij}) = -(n_{ij} - q_{ij}) \dots\dots\dots(2)$$

$$c_{ij} + l_{ij} + o_{ij} = f_{ij} + l_{ij} + r_{ij} \Rightarrow (c_{ij} - f_{ij}) + (l_{ij} - r_{ij}) = -(o_{ij} - l_{ij}) \dots\dots\dots(3)$$

Thus, we can reduce the linear system to the following

$$(a_{11} - d_{11})v_1 + (b_{11} - e_{11})v_2 + (c_{11} - f_{11})v_3 + (a_{12} - d_{12})v_4 + (b_{12} - e_{12})v_5 + (c_{12} - f_{12})v_6 = 0$$

$$(g_{11} - j_{11})v_1 + (h_{11} - k_{11})v_2 + (i_{11} - l_{11})v_3 + (g_{12} - j_{12})v_4 + (h_{12} - k_{12})v_5 + (i_{12} - l_{12})v_6 = 0$$

$$(a_{21} - d_{21})v_1 + (b_{21} - e_{21})v_2 + (c_{21} - f_{21})v_3 + (a_{22} - d_{22})v_4 + (b_{22} - e_{22})v_5 + (c_{22} - f_{22})v_6 = 0$$

$$(g_{21} - j_{21})v_1 + (h_{21} - k_{21})v_2 + (i_{21} - l_{21})v_3 + (g_{22} - j_{22})v_4 + (h_{22} - k_{22})v_5 + (i_{22} - l_{22})v_6 = 0$$

We can verify using the computer that the coefficient matrix of this system has in general the rank four. Hence, we deduce that v_1, v_2, v_3, v_4 depends on v_5 and v_6 .

By letting v_5 and v_6 take the values 0 and 1 we obtain two linearly independent vectors in the nullspace. These two vectors do not possess the property that the first six elements are the opposite of the last six elements. Hence, they are independent of the vector $(1, 1, 1, 1, 1, -1, -1, -1, -1, -1)^T$. \square

Remark: We did not here make use of the relation $B_{22} + B_{11} = B_{12} + B_{21}$. It actually does not affect the proof.

For example, the following square is a compound 12×12 magic square

-51	39	26	0	9	13	6	17	15	-6	0	4
54	-10	-2	-5	4	-5	20	5	2	0	9	0
-5	1	2	3	17	18	-24	6	7	8	19	20
12	3	-1	63	-27	-14	18	12	8	6	-5	-3
17	8	17	-42	22	14	12	3	12	-8	7	10
9	-5	-6	17	11	10	4	-7	-8	36	6	5
2	53	45	-131	33	34	59	31	34	-	24	25
-10	0	10	11	12	13	-44	15	14	16	17	18
-89	21	22	23	29	30	-108	26	27	28	31	32
143	-21	-22	10	-41	-33	149	-12	-13	-47	-19	-22
1	0	-1	22	12	2	-4	-5	-6	56	-3	-2
-11	-17	-18	101	-9	-10	-16	-19	-20	120	-14	-15

Using the computer we can verify that its nullity is 3. In other words, the constructed subspace is the nullspace itself.

We can generalize the previous result for an arbitrary number of squares involved in the compound square.

Theorem 1.3: Let A_{ij} be distinct pandiagonal magic square with magic constant $2s$ having the structure

$$A_{ij} = \begin{bmatrix} a_{ij} & b_{ij} & c_{ij} & d_{ij} \\ e_{ij} & f_{ij} & g_{ij} & h_{ij} \\ s - c_{ij} & s - d_{ij} & s - a_{ij} & s - b_{ij} \\ s - g_{ij} & s - h_{ij} & s - e_{ij} & s - f_{ij} \end{bmatrix}$$

such that $A_{ij} = A_{1j} + A_{i1} - A_{11}$ for $i, j = 1, \dots, n$. Assume that $(a_{11} + c_{12} - c_{11} - a_{12}) \neq 0$. Then, the following $4n \times 4n$ matrix

$$\begin{bmatrix} A_{11} & A_{12} & A_{13} & \dots & A_{1n} \\ A_{21} & A_{22} & A_{23} & \dots & A_{2n} \\ A_{31} & A_{32} & A_{33} & \dots & A_{3n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ A_{n1} & A_{n2} & A_{n3} & \dots & A_{nn} \end{bmatrix}$$

possess a $2n-3$ dimensional subspace of its nullspace, which is generated by the vectors

$$\begin{pmatrix} b_{11} - d_{11} - b_{12} + d_{12} \\ -(a_{11} + c_{12} - c_{11} - a_{12}) \\ -(b_{11} - d_{11} - b_{12} + d_{12}) \\ (a_{11} + c_{12} - c_{11} - a_{12}) \\ -(b_{11} - d_{11} - b_{12} + d_{12}) \\ a_{11} + c_{12} - c_{11} - a_{12} \\ (b_{11} - d_{11} - b_{12} + d_{12}) \\ -(a_{11} + c_{12} - c_{11} - a_{12}) \\ [0] \\ \vdots \\ [0] \end{pmatrix}$$

and

$$\begin{pmatrix}
 a_{12} + c_{13} - c_{12} - a_{13} \\
 0 \\
 -(a_{12} + c_{13} - c_{12} - a_{13}) \\
 0 \\
 -(a_{11} - c_{11} + c_{13} - a_{13}) \\
 0 \\
 a_{11} - c_{11} + c_{13} - a_{13} \\
 0 \\
 a_{11} + c_{12} - c_{11} - a_{12} \\
 0 \\
 -(a_{11} + c_{12} - c_{11} - a_{12}) \\
 0 \\
 [0] \\
 \vdots \\
 [0]
 \end{pmatrix}, \quad
 \begin{pmatrix}
 b_{11} - d_{11} - b_{13} + d_{13} \\
 -(a_{11} + c_{12} - c_{11} - a_{12}) \\
 -(b_{11} - d_{11} - b_{13} + d_{13}) \\
 a_{11} + c_{12} - c_{11} - a_{12} \\
 -(b_{11} - d_{11} - b_{13} + d_{13}) \\
 0 \\
 b_{11} - d_{11} - b_{13} + d_{13} \\
 0 \\
 0 \\
 [0] \\
 \vdots \\
 [0] \\
 \vdots \\
 [0]
 \end{pmatrix}, \quad
 \begin{pmatrix}
 a_{12} + c_{1n} - c_{12} - a_{1n} \\
 0 \\
 -(a_{12} + c_{1n} - c_{12} - a_{1n}) \\
 0 \\
 -(a_{11} - c_{11} + c_{1n} - a_{1n}) \\
 0 \\
 a_{11} - c_{11} + c_{1n} - a_{1n} \\
 0 \\
 a_{11} + c_{12} - c_{11} - a_{12} \\
 0 \\
 -(a_{11} + c_{12} - c_{11} - a_{12}) \\
 0 \\
 -(a_{11} + c_{12} - c_{11} - a_{12}) \\
 0
 \end{pmatrix}, \quad
 \begin{pmatrix}
 b_{11} - d_{11} - b_{1n} + d_{1n} \\
 -(a_{11} + c_{12} - c_{11} - a_{12}) \\
 -(b_{11} - d_{11} - b_{1n} + d_{1n}) \\
 a_{11} + c_{12} - c_{11} - a_{12} \\
 -(b_{11} - d_{11} - b_{1n} + d_{1n}) \\
 0 \\
 b_{11} - d_{11} - b_{1n} + d_{1n} \\
 0 \\
 [0] \\
 \vdots \\
 [0] \\
 0 \\
 a_{11} + c_{12} - c_{11} - a_{12} \\
 0 \\
 -(a_{11} + c_{12} - c_{11} - a_{12})
 \end{pmatrix}$$

Proof: We will check first that these vectors belong to the nullspace of the matrix. When we multiply the first vector with the matrix, we obtain a vector having in the first row

$$\begin{aligned}
 & (a_{11} - c_{11})(b_{11} - d_{11} - b_{12} + d_{12}) + (b_{11} - d_{11})(a_{11} - c_{11} - a_{12} + c_{12}) - (a_{12} - c_{12})(b_{11} - d_{11} - b_{12} + d_{12}) - \\
 & (b_{12} - d_{12})(a_{11} - c_{11} - a_{12} + c_{12}) = \\
 & (b_{11} - d_{11} - b_{12} + d_{12})[(a_{11} - c_{11}) - (a_{12} - c_{12})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{11} - d_{11}) - (b_{12} - d_{12})]\} = 0
 \end{aligned}$$

Since we know that

$$\begin{aligned}
 (a_{11} - c_{11}) &= -(e_{11} - g_{11}) \\
 (b_{11} - d_{11}) &= -(f_{11} - h_{11})
 \end{aligned}$$

we obtain zero in the second row of the vector. Since the third and fourth rows of the squares are complementary to the first two rows, we deduce that the third and fourth rows of the vector are also zero.

Now, the fifth entry of the vector is

$$\begin{aligned}
 & (a_{21} - c_{21})(b_{11} - d_{11} - b_{12} + d_{12}) + (b_{21} - d_{21})(a_{11} - c_{11} - a_{12} + c_{12}) - \\
 & (a_{22} - c_{22})(b_{11} - d_{11} - b_{12} + d_{12}) - (b_{22} - d_{22})(a_{11} - c_{11} - a_{12} + c_{12}) = \\
 & (b_{11} - d_{11} - b_{12} + d_{12})[(a_{21} - c_{21}) - (a_{22} - c_{22})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{21} - d_{21}) - (b_{22} - d_{22})]\}
 \end{aligned}$$

We use the following relations according to our assumption

$$\begin{aligned}a_{22} &= a_{12} + a_{21} - a_{11} \\b_{22} &= b_{12} + b_{21} - b_{11} \\c_{22} &= c_{12} + c_{21} - c_{11} \\d_{22} &= d_{12} + d_{21} - d_{11}\end{aligned}$$

and obtain

$$\begin{aligned}(b_{11} - d_{11} - b_{12} + d_{12})[(a_{21} - c_{21}) - (a_{12} + a_{21} - a_{11} - c_{12} - c_{21} + c_{11})] \\- \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{21} - d_{21}) - (b_{12} + b_{21} - b_{11} - d_{12} - d_{21} + d_{11})]\} \\= (b_{11} - d_{11} - b_{12} + d_{12})[-(a_{12} - a_{11} - c_{12} + c_{11})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[-(b_{12} - b_{11} - d_{12} + d_{11})]\} = 0\end{aligned}$$

We continue checking all rows until we reach the last entry, which is

$$\begin{aligned}(a_{n1} - c_{n1})(b_{11} - d_{11} - b_{12} + d_{12}) + (b_{n1} - d_{n1})(a_{11} - c_{11} - a_{12} + c_{12}) - \\(a_{n2} - c_{n2})(b_{11} - d_{11} - b_{12} + d_{12}) - (b_{n2} - d_{n2})(a_{11} - c_{11} - a_{12} + c_{12}) = \\(b_{11} - d_{11} - b_{12} + d_{12})[(a_{n1} - c_{n1}) - (a_{n2} - c_{n2})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{n1} - d_{n1}) - (b_{n2} - d_{n2})]\}\end{aligned}$$

We use

$$\begin{aligned}a_{n2} &= a_{12} + a_{n1} - a_{11} \\b_{n2} &= b_{12} + b_{n1} - b_{11} \\c_{n2} &= c_{12} + c_{n1} - c_{11} \\d_{n2} &= d_{12} + d_{n1} - d_{11}\end{aligned}$$

in order to obtain this value of the entry

$$\begin{aligned}(b_{11} - d_{11} - b_{12} + d_{12})[(a_{n1} - c_{n1}) - (a_{12} + a_{n1} - a_{11} - c_{12} - c_{n1} + c_{11})] \\- \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{n1} - d_{n1}) - (b_{12} + b_{n1} - b_{11} - d_{12} - d_{n1} + d_{11})]\} \\= (b_{11} - d_{11} - b_{12} + d_{12})[-(a_{12} - a_{11} - c_{12} + c_{11})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[-(b_{12} - b_{11} - d_{12} + d_{11})]\} = 0\end{aligned}$$

Hence, we finished checking the first vector.

Now, we turn our attention to the second vector. When we multiply the matrix with it, we obtain in the first entry

$$\begin{aligned}(a_{11} - c_{11})(a_{12} - c_{12} - a_{13} + c_{13}) - (a_{12} - c_{12})(a_{11} - c_{11} - a_{13} + c_{13}) + (a_{13} - c_{13})(a_{11} - c_{11} - a_{12} + c_{12}) = \\(a_{11} - c_{11})[(a_{12} - c_{12}) - (a_{13} - c_{13})] - (a_{12} - c_{12})[(a_{11} - c_{11}) - (a_{13} - c_{13})] + (a_{13} - c_{13})[(a_{11} - c_{11}) - (a_{12} - c_{12})] \\= (a_{11} - c_{11})(a_{12} - c_{12}) - (a_{11} - c_{11})(a_{13} - c_{13}) - (a_{12} - c_{12})(a_{11} - c_{11}) + (a_{12} - c_{12})(a_{13} - c_{13}) + (a_{13} - c_{13})(a_{11} - c_{11}) \\- (a_{13} - c_{13})(a_{12} - c_{12}) = 0\end{aligned}$$

Using the relations

$$\begin{aligned}(a_{11} - c_{11}) &= -(e_{11} - g_{11}) \\(b_{11} - d_{11}) &= -(f_{11} - h_{11})\end{aligned}$$

we deduce that the second entry is also zero. In a similar manner we can deal with the third and fourth entries. The fifth entry will be

$$(a_{21} - c_{21})(a_{12} - c_{12} - a_{13} + c_{13}) - (a_{22} - c_{22})(a_{11} - c_{11} - a_{13} + c_{13}) + (a_{23} - c_{23})(a_{11} - c_{11} - a_{12} + c_{12})$$

We use the relations

$$\begin{aligned} a_{22} &= a_{12} + a_{21} - a_{11} \\ c_{22} &= c_{12} + c_{21} - c_{11} \\ a_{23} &= a_{13} + a_{21} - a_{11} \\ c_{23} &= c_{13} + c_{21} - c_{11} \end{aligned}$$

to obtain for the fifth entry

$$\begin{aligned} &= (a_{21} - c_{21})[(a_{12} - c_{12}) - (a_{13} - c_{13})] - (a_{12} + a_{21} - a_{11} - c_{12} - c_{21} + c_{11})[(a_{11} - c_{11}) - (a_{13} - c_{13})] \\ &\quad + (a_{13} + a_{21} - a_{11} - c_{13} - c_{21} + c_{11})[(a_{11} - c_{11}) - (a_{12} - c_{12})] \\ &= (a_{21} - c_{21})(a_{12} - c_{12}) - (a_{21} - c_{21})(a_{13} - c_{13}) - (a_{21} - c_{21})(a_{11} - c_{11}) + (a_{21} - c_{21})(a_{13} - c_{13}) \\ &\quad - (a_{12} - c_{12})(a_{11} - c_{11}) + (a_{12} - c_{12})(a_{13} - c_{13}) + (a_{11} - c_{11})^2 + (a_{11} - c_{11})(a_{13} - c_{13}) + (a_{13} - c_{13})(a_{11} - c_{11}) \\ &\quad - (a_{13} - c_{13})(a_{12} - c_{12}) + (a_{21} - c_{21})(a_{11} - c_{11}) - (a_{21} - c_{21})(a_{12} - c_{12}) - (a_{11} - c_{11})^2 + (a_{12} - c_{12})(a_{11} - c_{11}) = 0 \end{aligned}$$

We continue checking the entries until we reach the last entry, which is

$$(a_{n1} - c_{n1})(a_{12} - c_{12} - a_{13} + c_{13}) - (a_{n2} - c_{n2})(a_{11} - c_{11} - a_{13} + c_{13}) + (a_{n3} - c_{n3})(a_{11} - c_{11} - a_{12} + c_{12})$$

Using the relations

$$\begin{aligned} a_{n2} &= a_{12} + a_{n1} - a_{11} \\ c_{n2} &= c_{12} + c_{n1} - c_{11} \\ a_{n3} &= a_{13} + a_{n1} - a_{11} \\ c_{n3} &= c_{13} + c_{n1} - c_{11} \end{aligned}$$

we get

$$\begin{aligned} &= (a_{n1} - c_{n1})[(a_{12} - c_{12}) - (a_{13} - c_{13})] - (a_{12} + a_{n1} - a_{11} - c_{12} - c_{n1} + c_{11})[(a_{11} - c_{11}) - (a_{13} - c_{13})] \\ &\quad + (a_{13} + a_{n1} - a_{11} - c_{13} - c_{n1} + c_{11})[(a_{11} - c_{11}) - (a_{12} - c_{12})] \\ &= (a_{n1} - c_{n1})(a_{12} - c_{12}) - (a_{n1} - c_{n1})(a_{13} - c_{13}) - (a_{n1} - c_{n1})(a_{11} - c_{11}) + (a_{n1} - c_{n1})(a_{13} - c_{13}) \\ &\quad - (a_{12} - c_{12})(a_{11} - c_{11}) + (a_{12} - c_{12})(a_{13} - c_{13}) + (a_{11} - c_{11})^2 + (a_{11} - c_{11})(a_{13} - c_{13}) + (a_{13} - c_{13})(a_{11} - c_{11}) \\ &\quad - (a_{13} - c_{13})(a_{12} - c_{12}) + (a_{n1} - c_{n1})(a_{11} - c_{11}) - (a_{n1} - c_{n1})(a_{12} - c_{12}) - (a_{11} - c_{11})^2 + (a_{12} - c_{12})(a_{11} - c_{11}) = 0 \end{aligned}$$

Hence, the second vector belongs to the nullspace of the $(4n \times 4n)$ -matrix.

Similarly, we can check that all the other vectors are included in the nullspace of the $(4n \times 4n)$ -matrix. We check the last vector (the $(2n - 3)$ -th vector) belongs to the nullspace of the $(4n \times 4n)$ -matrix. The first entry by matrix multiplication is

$$\begin{aligned} &(a_{11} - c_{11})(b_{11} - d_{11} - b_{1n} + d_{1n}) + (b_{11} - d_{11})(a_{11} - c_{11} - a_{12} + c_{12}) - \\ &(a_{12} - c_{12})(b_{11} - d_{11} - b_{1n} + d_{1n}) - (b_{1n} - d_{1n})(a_{11} - c_{11} - a_{12} + c_{12}) = \\ &(b_{11} - d_{11} - b_{1n} + d_{1n})[(a_{11} - c_{11}) - (a_{12} - c_{12})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{11} - d_{11}) - (b_{1n} - d_{1n})]\} = 0 \end{aligned}$$

As before we deuce also that the second, third and fourth entries are zero. The fifth entry is

$$\begin{aligned}
& (a_{21} - c_{21})(b_{11} - d_{11} - b_{1n} + d_{1n}) + (b_{21} - d_{21})(a_{11} - c_{11} - a_{12} + c_{12}) - \\
& (a_{22} - c_{22})(b_{11} - d_{11} - b_{1n} + d_{1n}) - (b_{2n} - d_{2n})(a_{11} - c_{11} - a_{12} + c_{12}) = \\
& (b_{11} - d_{11} - b_{1n} + d_{1n})[(a_{21} - c_{21}) - (a_{22} - c_{22})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{21} - d_{21}) - (b_{2n} - d_{2n})]\} \\
& = (b_{11} - d_{11} - b_{1n} + d_{1n})[(a_{21} - c_{21}) - (a_{12} + a_{21} - a_{11} - c_{12} - c_{21} + c_{11})]
\end{aligned}$$

We use the relations

$$\begin{aligned}
a_{22} &= a_{12} + a_{21} - a_{11} \\
b_{2n} &= b_{1n} + b_{21} - b_{11} \\
c_{22} &= c_{12} + c_{21} - c_{11} \\
d_{2n} &= d_{1n} + d_{21} - d_{11}
\end{aligned}$$

Therefore, this entry is

$$\begin{aligned}
& (b_{11} - d_{11} - b_{1n} + d_{1n})[(a_{21} - c_{21}) - (a_{12} + a_{21} - a_{11} - c_{12} - c_{21} + c_{11})] \\
& - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{21} - d_{21}) - (b_{1n} + b_{21} - b_{11} - d_{1n} - d_{21} + d_{11})]\} \\
& = (b_{11} - d_{11} - b_{1n} + d_{1n})[-(a_{12} - a_{11} - c_{12} + c_{11})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[-(b_{1n} - b_{11} - d_{1n} + d_{11})]\} = 0
\end{aligned}$$

When we reach the ($2n - 3$)-th entry, we find that it is

$$\begin{aligned}
& (a_{n1} - c_{n1})(b_{11} - d_{11} - b_{1n} + d_{1n}) + (b_{n1} - d_{n1})(a_{11} - c_{11} - a_{12} + c_{12}) - \\
& (a_{nn} - c_{nn})(b_{11} - d_{11} - b_{1n} + d_{1n}) - (b_{nn} - d_{nn})(a_{11} - c_{11} - a_{12} + c_{12}) = \\
& (b_{11} - d_{11} - b_{1n} + d_{1n})[(a_{n1} - c_{n1}) - (a_{nn} - c_{nn})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{n1} - d_{n1}) - (b_{nn} - d_{nn})]\}
\end{aligned}$$

We use the relations

$$\begin{aligned}
a_{nn} &= a_{1n} + a_{n1} - a_{11} \\
b_{nn} &= b_{1n} + b_{n1} - b_{11} \\
c_{nn} &= c_{1n} + c_{n1} - c_{11} \\
d_{nn} &= d_{1n} + d_{n1} - d_{11}
\end{aligned}$$

to prove that this entry is

$$\begin{aligned}
& (b_{11} - d_{11} - b_{1n} + d_{1n})[(a_{n1} - c_{n1}) - (a_{12} + a_{n1} - a_{11} - c_{12} - c_{n1} + c_{11})] \\
& - \{(a_{11} - c_{11} - a_{12} + c_{12})[(b_{n1} - d_{n1}) - (b_{1n} + b_{n1} - b_{11} - d_{1n} - d_{n1} + d_{11})]\} \\
& = (b_{11} - d_{11} - b_{1n} + d_{1n})[-(a_{12} - a_{11} - c_{12} + c_{11})] - \{(a_{11} - c_{11} - a_{12} + c_{12})[-(b_{1n} - b_{11} - d_{1n} + d_{11})]\} = 0
\end{aligned}$$

We prove now that the vectors are linearly independent. Let $k_1, k_2, k_3, \dots, k_{2n-4}, k_{2n-3} \in R$ such that

This leads us to

From the $(4n-2)$ -th row of this vector we obtain the equation

$$k_{2n-3}(a_{11} + c_{12} - c_{11} - a_{12}) = 0$$

According to our assumptions we must have $k_{2n-3} = 0$. Similarly, we obtain $k_{2n-4} = 0$ from the $(4n-3)$ -th row. We continue checking all the rows up to the tenth row, which looks like this

$$k_3(a_{11} + c_{12} - c_{11} - a_{12}) = 0$$

Hence, we conclude that $k_3 = 0$. From the ninth (res. eighth) row we obtain $k_2 = 0$ (res. $k_1 = 0$). Since all $k_1, k_2, k_3, \dots, k_{2n-4}, k_{2n-3}$ are zero, we are done. \square

We prove now a similar result to the previous theorem, where we replace the 4×4 square with a 6×6 one.

Theorem 2.4: Let B_{ij} be 6×6 panmagic squares with magic constant $3s$ such that

$$B_{ij} = B_{1j} + B_{i1} - B_{11} \text{ for all } i, j = 1, 2, 3, \dots, n.$$

Moreover, we assume that $(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2) \neq 0$. Then the following square

$$\begin{bmatrix} B_{11} & B_{12} & B_{13} & \dots & B_{1n} \\ B_{21} & B_{22} & B_{23} & \dots & B_{2n} \\ B_{31} & B_{32} & B_{33} & \dots & B_{3n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ B_{n1} & B_{n2} & B_{n3} & \dots & B_{nn} \end{bmatrix}$$

is a magic square with magic constant $3ns$. Further, it has a $(3n-5)$ -dimensional subspace of its nullspace. This space is generated by the vectors:

$$\left(\begin{array}{c} -[(b_1 - b_2)(i_1 - i_2) - (c_1 - c_2)(h_1 - h_2)] \\ [(a_1 - a_2)(i_1 - i_2) - (c_1 - c_2)(g_1 - g_2)] \\ -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] \\ [(b_1 - b_2)(i_1 - i_2) - (c_1 - c_2)(h_1 - h_2)] \\ -[(a_1 - a_2)(i_1 - i_2) - (c_1 - c_2)(g_1 - g_2)] \\ [(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] \\ [(b_1 - b_2)(i_1 - i_2) - (c_1 - c_2)(h_1 - h_2)] \\ -[(a_1 - a_2)(i_1 - i_2) - (c_1 - c_2)(g_1 - g_2)] \\ (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2) \\ -[(b_1 - b_2)(i_1 - i_2) - (c_1 - c_2)(h_1 - h_2)] \\ [(a_1 - a_2)(i_1 - i_2) - (c_1 - c_2)(g_1 - g_2)] \\ -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] \\ [0] \\ [0] \\ \vdots \\ [0] \end{array} \right)$$

$$\begin{pmatrix}
 (a_2 - a_3)(h_1 - h_2) - (b_1 - b_2)(g_2 - g_3) & & & \\
 a_1(g_2 - g_3) - a_2(g_1 - g_3) + a_3(g_1 - g_2) & \left(-[b_1(h_2 - h_3) - b_2(h_1 - h_3) + b_3(h_1 - h_2)] \right) & \left(-(b_1 - b_2)(i_1 - i_3) - (c_1 - c_3)(h_1 - h_2) \right) & \\
 0 & (a_1 - a_2)(h_2 - h_3) - (b_2 - b_3)(g_1 - g_2) & (a_1 - a_2)(i_1 - i_3) - (c_1 - c_3)(g_1 - g_2) & \\
 -[(a_2 - a_3)(h_1 - h_2) - (b_1 - b_2)(g_2 - g_3)] & 0 & -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & \\
 -[a_1(g_2 - g_3) - a_2(g_1 - g_3) + a_3(g_1 - g_2)] & [b_1(h_2 - h_3) - b_2(h_1 - h_3) + b_3(h_1 - h_2)] & [(b_1 - b_2)(i_1 - i_3) - (c_1 - c_3)(h_1 - h_2)] & \\
 0 & -[(a_1 - a_2)(h_2 - h_3) - (b_2 - b_3)(g_1 - g_2)] & -[(a_1 - a_2)(i_1 - i_3) - (c_1 - c_3)(g_1 - g_2)] & \\
 -(a_1 - a_3)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_3) & 0 & [(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & \\
 -a_1(g_2 - g_3) + a_2(g_1 - g_3) - a_3(g_1 - g_2) & b_1(h_2 - h_3) - b_2(h_1 - h_3) + b_3(h_1 - h_2) & (b_1 - b_2)(i_1 - i_3) - (c_1 - c_3)(h_1 - h_2) & \\
 0 & -[(a_1 - a_2)(h_1 - h_3) - (b_1 - b_3)(g_1 - g_2)] & -[(a_1 - a_2)(i_1 - i_3) - (c_1 - c_3)(g_1 - g_2)] & \\
 (a_1 - a_3)(h_1 - h_2) + (b_1 - b_2)(g_1 - g_3) & 0 & 0 & \\
 a_1(g_2 - g_3) - a_2(g_1 - g_3) + a_3(g_1 - g_2) & \left[(a_1 - a_2)(h_1 - h_3) - (b_1 - b_3)(g_1 - g_2) \right] & \left[(a_1 - a_2)(i_1 - i_3) - (c_1 - c_3)(h_1 - h_2) \right] & \\
 0 & 0 & 0 & \\
 [0] & [0] & [0] & \\
 (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2) & 0 & 0 & \\
 0 & (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2) & 0 & \\
 0 & 0 & 0 & \\
 -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & 0 & 0 & \\
 0 & -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & 0 & \\
 0 & 0 & -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & \\
 [0] & [0] & [0] & \\
 \vdots & \vdots & \vdots & \\
 [0] & [0] & [0] &
 \end{pmatrix},$$

$$\begin{pmatrix}
 (a_2 - a_n)(h_1 - h_2) - (b_1 - b_2)(g_2 - g_n) & & & \\
 a_1(g_2 - g_n) - a_2(g_1 - g_n) + a_n(g_1 - g_2) & \left(-[b_1(h_2 - h_n) - b_2(h_1 - h_n) + b_n(h_1 - h_2)] \right) & \left(-(b_1 - b_2)(i_1 - i_n) - (c_1 - c_n)(h_1 - h_2) \right) & \\
 0 & (a_1 - a_2)(h_2 - h_n) - (b_2 - b_n)(g_1 - g_2) & (a_1 - a_2)(i_1 - i_n) - (c_1 - c_n)(g_1 - g_2) & \\
 -[(a_2 - a_n)(h_1 - h_2) - (b_1 - b_2)(g_2 - g_n)] & 0 & -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & \\
 -[a_1(g_2 - g_n) - a_2(g_1 - g_n) + a_n(g_1 - g_2)] & [b_1(h_2 - h_n) - b_2(h_1 - h_n) + b_n(h_1 - h_2)] & [(b_1 - b_2)(i_1 - i_n) - (c_1 - c_n)(h_1 - h_2)] & \\
 0 & -[(a_1 - a_2)(h_2 - h_n) - (b_2 - b_n)(g_1 - g_2)] & -[(a_1 - a_2)(i_1 - i_n) - (c_1 - c_n)(g_1 - g_2)] & \\
 -(a_1 - a_n)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_n) & 0 & [(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & \\
 -a_1(g_2 - g_n) + a_2(g_1 - g_n) - a_n(g_1 - g_2) & b_1(h_2 - h_n) - b_2(h_1 - h_n) + b_n(h_1 - h_2) & (b_1 - b_2)(i_1 - i_n) - (c_1 - c_n)(h_1 - h_2) & \\
 0 & -[(a_1 - a_2)(h_1 - h_n) - (b_1 - b_n)(g_1 - g_2)] & -[(a_1 - a_2)(i_1 - i_n) - (c_1 - c_n)(g_1 - g_2)] & \\
 (a_1 - a_n)(h_1 - h_2) + (b_1 - b_2)(g_1 - g_n) & 0 & 0 & \\
 , a_1(g_2 - g_n) - a_2(g_1 - g_n) + a_n(g_1 - g_2) & \left[(a_1 - a_2)(h_1 - h_n) - (b_1 - b_n)(g_1 - g_2) \right] & \left[(a_1 - a_2)(i_1 - i_n) - (c_1 - c_n)(g_1 - g_2) \right] & \\
 0 & 0 & 0 & \\
 [0] & [0] & [0] & \\
 \vdots & \vdots & \vdots & \\
 [0] & [0] & [0] & \\
 (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2) & 0 & 0 & \\
 0 & (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2) & 0 & \\
 0 & 0 & 0 & \\
 -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & 0 & 0 & \\
 0 & -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & 0 & \\
 0 & 0 & -[(a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)] & \\
 \end{pmatrix},$$

Proof: It is easy to prove that it is a magic square. We look now for vectors in R^{6n} having the following structure

$$(v_1, v_2, v_3, -v_1 - v_2 - v_3, \dots, v_{3n-2}, v_{3n-1}, v_{3n}, -v_{3n-2}, -v_{3n-1}, -v_{3n})'$$

where $v_1, v_2, v_3, v_4, v_5, v_6, v_7, v_8, v_9, \dots, v_{3n-2}, v_{3n-1}, v_{3n}$ are scalars, which can be determined by solving

$$\begin{bmatrix} B_{11} & B_{12} & B_{13} & \dots & B_{1n} \\ B_{21} & B_{22} & B_{23} & \dots & B_{2n} \\ B_{31} & B_{32} & B_{33} & \dots & B_{3n} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ B_{n1} & B_{n2} & B_{n3} & \dots & B_{nn} \end{bmatrix} \begin{pmatrix} v_1 \\ v_2 \\ v_3 \\ \vdots \\ v_{3n} \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

Using the structure of the vector we can eliminate the terms including s . Moreover, the resulting linear system has repeated equations. In fact, the number of equations, which we have to solve, is $3n$. They are

$$\begin{aligned} (a_{11} - d_{11})v_1 + (b_{11} - e_{11})v_2 + (c_{11} - f_{11})v_3 + (a_{12} - d_{12})v_4 + (b_{12} - e_{12})v_5 + (c_{12} - f_{12})v_6 + (a_{13} - d_{13})v_7 + (b_{13} - e_{13})v_8 + (c_{13} - f_{13})v_9 + \dots + (a_{1n} - d_{1n})v_{3n-2} + (b_{1n} - e_{1n})v_{3n-1} + (c_{1n} - f_{1n})v_{3n} &= 0 \\ (g_{11} - j_{11})v_1 + (h_{11} - k_{11})v_2 + (i_{11} - l_{11})v_3 + (g_{12} - j_{12})v_4 + (h_{12} - k_{12})v_5 + (i_{12} - l_{12})v_6 + (g_{13} - j_{13})v_7 + (h_{13} - k_{13})v_8 + (i_{13} - l_{13})v_9 + \dots + (g_{1n} - j_{1n})v_{3n-2} + (h_{1n} - k_{1n})v_{3n-1} + (i_{1n} - l_{1n})v_{3n} &= 0 \\ (m_{11} - p_{11})v_1 + (n_{11} - q_{11})v_2 + (o_{11} - r_{11})v_3 + (m_{12} - p_{12})v_4 + (n_{12} - q_{12})v_5 + (o_{12} - r_{12})v_6 + (m_{13} - p_{13})v_7 + (n_{13} - q_{13})v_8 + (o_{13} - r_{13})v_9 + \dots + (m_{1n} - p_{1n})v_{3n-2} + (n_{1n} - q_{1n})v_{3n-1} + (o_{1n} - r_{1n})v_{3n} &= 0 \\ (a_{21} - d_{21})v_1 + (b_{21} - e_{21})v_2 + (c_{21} - f_{21})v_3 + (a_{22} - d_{22})v_4 + (b_{22} - e_{22})v_5 + (c_{22} - f_{22})v_6 + (a_{23} - d_{23})v_7 + (b_{23} - e_{23})v_8 + (c_{23} - f_{23})v_9 + \dots + (a_{2n} - d_{2n})v_{3n-2} + (b_{2n} - e_{2n})v_{3n-1} + (c_{2n} - f_{2n})v_{3n} &= 0 \\ (g_{21} - j_{21})v_1 + (h_{21} - k_{21})v_2 + (i_{21} - l_{21})v_3 + (g_{22} - j_{22})v_4 + (h_{22} - k_{22})v_5 + (i_{22} - l_{22})v_6 + (g_{23} - j_{23})v_7 + (h_{23} - k_{23})v_8 + (i_{23} - l_{23})v_9 + \dots + (g_{2n} - j_{2n})v_{3n-2} + (h_{2n} - k_{2n})v_{3n-1} + (i_{2n} - l_{2n})v_{3n} &= 0 \\ (m_{21} - p_{21})v_1 + (n_{21} - q_{21})v_2 + (o_{21} - r_{21})v_3 + (m_{22} - p_{22})v_4 + (n_{22} - q_{22})v_5 + (o_{22} - r_{22})v_6 + (m_{23} - p_{23})v_7 + (n_{23} - q_{23})v_8 + (o_{23} - r_{23})v_9 + \dots + (m_{2n} - p_{2n})v_{3n-2} + (n_{2n} - q_{2n})v_{3n-1} + (o_{2n} - r_{2n})v_{3n} &= 0 \\ (a_{31} - d_{31})v_1 + (b_{31} - e_{31})v_2 + (c_{31} - f_{31})v_3 + (a_{32} - d_{32})v_4 + (b_{32} - e_{32})v_5 + (c_{32} - f_{32})v_6 + (a_{33} - d_{33})v_7 + (b_{33} - e_{33})v_8 + (c_{33} - f_{33})v_9 + \dots + (a_{3n} - d_{3n})v_{3n-2} + (b_{3n} - e_{3n})v_{3n-1} + (c_{3n} - f_{3n})v_{3n} &= 0 \\ (g_{31} - j_{31})v_1 + (h_{31} - k_{31})v_2 + (i_{31} - l_{31})v_3 + (g_{32} - j_{32})v_4 + (h_{32} - k_{32})v_5 + (i_{32} - l_{32})v_6 + (g_{33} - j_{33})v_7 + (h_{33} - k_{33})v_8 + (i_{33} - l_{33})v_9 + \dots + (g_{3n} - j_{3n})v_{3n-2} + (h_{3n} - k_{3n})v_{3n-1} + (i_{3n} - l_{3n})v_{3n} &= 0 \\ (m_{31} - p_{31})v_1 + (n_{31} - q_{31})v_2 + (o_{31} - r_{31})v_3 + (m_{32} - p_{32})v_4 + (n_{32} - q_{32})v_5 + (o_{32} - r_{32})v_6 + (m_{33} - p_{33})v_7 + (n_{33} - q_{33})v_8 + (o_{33} - r_{33})v_9 + \dots + (m_{3n} - p_{3n})v_{3n-2} + (n_{3n} - q_{3n})v_{3n-1} + (o_{3n} - r_{3n})v_{3n} &= 0 \end{aligned}$$

⋮

$$\begin{aligned}
 & (a_{n1} - d_{n1})v_1 + (b_{n1} - e_{n1})v_2 + (c_{n1} - f_{n1})v_3 + (a_{n2} - d_{n2})v_4 + (b_{n2} - e_{n2})v_5 + (c_{n2} - f_{n2})v_6 + (a_{n3} - d_{n3})v_7 + \\
 & (b_{n3} - e_{n3})v_8 + (c_{n3} - f_{n3})v_9 + \dots + (a_{nn} - d_{nn})v_{3n-2} + (b_{nn} - e_{nn})v_{3n-1} + (c_{nn} - f_{nn})v_{3n} = 0 \\
 & (g_{n1} - j_{n1})v_1 + (h_{n1} - k_{n1})v_2 + (i_{n1} - l_{n1})v_3 + (g_{n2} - j_{n2})v_4 + (h_{n2} - k_{n2})v_5 + (i_{n2} - l_{n2})v_6 + (g_{n3} - j_{n3})v_7 + \\
 & (h_{n3} - k_{n3})v_8 + (i_{n3} - l_{n3})v_9 + \dots + (g_{nn} - j_{nn})v_{3n-2} + (h_{nn} - k_{nn})v_{3n-1} + (i_{nn} - l_{nn})v_{3n} = 0 \\
 & (m_{n1} - p_{n1})v_1 + (n_{n1} - q_{n1})v_2 + (o_{n1} - r_{n1})v_3 + (m_{n2} - p_{n2})v_4 + (n_{n2} - q_{n2})v_5 + (o_{n2} - r_{n2})v_6 + (m_{n3} - p_{n3})v_7 + \\
 & (n_{n3} - q_{n3})v_8 + (o_{n3} - r_{n3})v_9 + \dots + (m_{nn} - p_{nn})v_{3n-2} + (n_{nn} - q_{nn})v_{3n-1} + (o_{nn} - r_{nn})v_{3n} = 0
 \end{aligned}$$

We know that the following relation holds for panmagic squares

$$a_{ij} + g_{ij} + m_{ij} = d_{ij} + j_{ij} + p_{ij} \Rightarrow (a_{ij} - d_{ij}) + (g_{ij} - j_{ij}) = -(m_{ij} - p_{ij}) \quad \dots\dots\dots(1)$$

$$b_{ij} + h_{ij} + n_{ij} = e_{ij} + k_{ij} + q_{ij} \Rightarrow (b_{ij} - e_{ij}) + (h_{ij} - k_{ij}) = -(n_{ij} - q_{ij}) \quad \dots\dots\dots(2)$$

$$c_{ij} + l_{ij} + o_{ij} = f_{ij} + i_{ij} + r_{ij} \Rightarrow (c_{ij} - f_{ij}) + (i_{ij} - l_{ij}) = -(o_{ij} - r_{ij}) \quad \dots\dots\dots(3)$$

Hence, we can reduce the previous system to the following system consisting of $2n$ equations

$$\begin{aligned}
 & (a_{11} - d_{11})v_1 + (b_{11} - e_{11})v_2 + (c_{11} - f_{11})v_3 + (a_{12} - d_{12})v_4 + (b_{12} - e_{12})v_5 + (c_{12} - f_{12})v_6 + (a_{13} - d_{13})v_7 + \\
 & (b_{13} - e_{13})v_8 + (c_{13} - f_{13})v_9 + \dots + (a_{1n} - d_{1n})v_{3n-2} + (b_{1n} - e_{1n})v_{3n-1} + (c_{1n} - f_{1n})v_{3n} = 0 \\
 & (g_{11} - j_{11})v_1 + (h_{11} - k_{11})v_2 + (i_{11} - l_{11})v_3 + (g_{12} - j_{12})v_4 + (h_{12} - k_{12})v_5 + (i_{12} - l_{12})v_6 + (g_{13} - j_{13})v_7 + \\
 & (h_{13} - k_{13})v_8 + (i_{13} - l_{13})v_9 + \dots + (g_{1n} - j_{1n})v_{3n-2} + (h_{1n} - k_{1n})v_{3n-1} + (i_{1n} - l_{1n})v_{3n} = 0 \\
 & (a_{21} - d_{21})v_1 + (b_{21} - e_{21})v_2 + (c_{21} - f_{21})v_3 + (a_{22} - d_{22})v_4 + (b_{22} - e_{22})v_5 + (c_{22} - f_{22})v_6 + (a_{23} - d_{23})v_7 + \\
 & (b_{23} - e_{23})v_8 + (c_{23} - f_{23})v_9 + \dots + (a_{2n} - d_{2n})v_{3n-2} + (b_{2n} - e_{2n})v_{3n-1} + (c_{2n} - f_{2n})v_{3n} = 0 \\
 & (g_{21} - j_{21})v_1 + (h_{21} - k_{21})v_2 + (i_{21} - l_{21})v_3 + (g_{22} - j_{22})v_4 + (h_{22} - k_{22})v_5 + (i_{22} - l_{22})v_6 + (g_{23} - j_{23})v_7 + \\
 & (h_{23} - k_{23})v_8 + (i_{23} - l_{23})v_9 + \dots + (g_{2n} - j_{2n})v_{3n-2} + (h_{2n} - k_{2n})v_{3n-1} + (i_{2n} - l_{2n})v_{3n} = 0 \\
 & (a_{31} - d_{31})v_1 + (b_{31} - e_{31})v_2 + (c_{31} - f_{31})v_3 + (a_{32} - d_{32})v_4 + (b_{32} - e_{32})v_5 + (c_{32} - f_{32})v_6 + (a_{33} - d_{33})v_7 + \\
 & (b_{33} - e_{33})v_8 + (c_{33} - f_{33})v_9 + \dots + (a_{3n} - d_{3n})v_{3n-2} + (b_{3n} - e_{3n})v_{3n-1} + (c_{3n} - f_{3n})v_{3n} = 0 \\
 & (g_{31} - j_{31})v_1 + (h_{31} - k_{31})v_2 + (i_{31} - l_{31})v_3 + (g_{32} - j_{32})v_4 + (h_{32} - k_{32})v_5 + (i_{32} - l_{32})v_6 + (g_{33} - j_{33})v_7 + \\
 & (h_{33} - k_{33})v_8 + (i_{33} - l_{33})v_9 + \dots + (g_{3n} - j_{3n})v_{3n-2} + (h_{3n} - k_{3n})v_{3n-1} + (i_{3n} - l_{3n})v_{3n} = 0
 \end{aligned}$$

⋮

$$\begin{aligned}
 & (a_{n1} - d_{n1})v_1 + (b_{n1} - e_{n1})v_2 + (c_{n1} - f_{n1})v_3 + (a_{n2} - d_{n2})v_4 + (b_{n2} - e_{n2})v_5 + (c_{n2} - f_{n2})v_6 + (a_{n3} - d_{n3})v_7 + \\
 & (b_{n3} - e_{n3})v_8 + (c_{n3} - f_{n3})v_9 + \dots + (a_{nn} - d_{nn})v_{3n-2} + (b_{nn} - e_{nn})v_{3n-1} + (c_{nn} - f_{nn})v_{3n} = 0 \\
 & (g_{n1} - j_{n1})v_1 + (h_{n1} - k_{n1})v_2 + (i_{n1} - l_{n1})v_3 + (g_{n2} - j_{n2})v_4 + (h_{n2} - k_{n2})v_5 + (i_{n2} - l_{n2})v_6 + (g_{n3} - j_{n3})v_7 + \\
 & (h_{n3} - k_{n3})v_8 + (i_{n3} - l_{n3})v_9 + \dots + (g_{nn} - j_{nn})v_{3n-2} + (h_{nn} - k_{nn})v_{3n-1} + (i_{nn} - l_{nn})v_{3n} = 0
 \end{aligned}$$

In order to simplify the calculations we use the following notations

$$a_1 = a_{11} - d_{11}, b_1 = b_{11} - e_{11}, c_1 = c_{11} - f_{11}, \dots, a_n = a_{1n} - d_{1n}, b_n = b_{1n} - e_{1n}, c_n = c_{1n} - f_{1n}$$

$$g_1 = g_{11} - j_{11}, h_1 = h_{11} - k_{11}, i_1 = i_{11} - l_{11}, \dots, g_n = g_{1n} - j_{1n}, h_n = h_{1n} - k_{1n}, i_n = i_{1n} - l_{1n}$$

Using the computer we deduce from the last system that

$$\begin{aligned}
 v_1 &= -\frac{(b_1-b_2)(i_1-i_2)-(c_1-c_2)(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_6 - \frac{-(a_2-a_3)(h_1-h_2)+(b_1-b_2)(g_2-g_3)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_7 \\
 &\quad - \frac{b_1(h_2-h_3)-b_2(h_1-h_3)+b_3(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_8 - \frac{(b_1-b_2)(i_1-i_3)-(c_1-c_3)(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_9 - \dots \\
 &\quad - \frac{-(a_2-a_n)(h_1-h_2)+(b_1-b_2)(g_2-g_n)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-2} - \frac{b_1(h_2-h_n)-b_2(h_1-h_n)+b_n(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-1} \\
 &\quad - \frac{(b_1-b_2)(i_1-i_n)-(c_1-c_n)(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n} \\
 \\
 v_2 &= \frac{(a_1-a_2)(i_1-i_2)-(c_1-c_2)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_6 + \frac{a_1(g_2-g_3)-a_2(g_1-g_3)+a_3(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_7 \\
 &\quad + \frac{(a_1-a_2)(h_2-h_3)-(b_2-b_3)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_8 + \frac{(a_1-a_2)(i_1-i_3)-(c_1-c_3)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_9 + \dots \\
 &\quad + \frac{a_1(g_2-g_n)-a_2(g_1-g_n)+a_n(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-2} + \frac{(a_1-a_2)(h_2-h_n)-(b_2-b_n)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-1} \\
 &\quad + \frac{(a_1-a_2)(i_1-i_n)-(c_1-c_n)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n} - \frac{(a_1-a_2)(h_1-h_n)-(b_1-b_n)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-1} \\
 &\quad - \frac{(a_1-a_2)(i_1-i_n)-(c_1-c_n)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n} \\
 \\
 v_3 &= -v_6 - v_9 - v_{12} - \dots - v_{3n} \\
 v_4 &= \frac{(b_1-b_2)(i_1-i_2)-(c_1-c_2)(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_6 - \frac{(a_1-a_3)(h_1-h_2)+(b_1-b_2)(g_1-g_3)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_7 + \frac{b_1(h_2-h_3)-b_2(h_1-h_3)+b_3(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_8 \\
 &\quad + \frac{(b_1-b_2)(i_1-i_3)-(c_1-c_3)(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_9 - \dots - \frac{(a_1-a_n)(h_1-h_2)+(b_1-b_2)(g_1-g_n)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-2} \\
 &\quad + \frac{b_1(h_2-h_n)-b_2(h_1-h_n)+b_n(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-1} + \frac{(b_1-b_2)(i_1-i_n)-(c_1-c_n)(h_1-h_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n} \\
 \\
 v_5 &= -\frac{(a_1-a_2)(i_1-i_2)-(c_1-c_2)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_6 - \frac{a_1(g_2-g_3)-a_2(g_1-g_3)+a_3(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_7 - \frac{(a_1-a_2)(h_1-h_3)-(b_1-b_3)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_8 \\
 &\quad - \frac{(a_1-a_2)(i_1-i_3)-(c_1-c_3)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_9 - \dots - \frac{a_1(g_2-g_n)-a_2(g_1-g_n)+a_n(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-2} \\
 &\quad - \frac{(a_1-a_2)(h_1-h_n)-(b_1-b_n)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n-1} - \frac{(a_1-a_2)(i_1-i_n)-(c_1-c_n)(g_1-g_2)}{(a_1-a_2)(h_1-h_2)-(b_1-b_2)(g_1-g_2)}v_{3n}
 \end{aligned}$$

If we set $v_6 = (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)$ and $v_7, v_8, v_9, \dots, v_{3n-2}, v_{3n-1}, v_{3n}$ to zero, then we obtain the first vector. If we set $v_7 = (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)$ and $v_6, v_8, v_9, \dots, v_{3n-2}, v_{3n-1}, v_{3n}$ to zero, then we obtain the second vector. We continue in this manner until we set

$$v_{3n} = (a_1 - a_2)(h_1 - h_2) - (b_1 - b_2)(g_1 - g_2)$$

and $v_6, v_7, v_8, v_9, \dots, v_{3n-2}, v_{3n-1}$ to zero, in order to obtain the last vector. It is clear from the way of their generation that the vectors are linearly independent. It is

also easy to multiply the square with each vector to verify that the result is the zero. \square

For example, we consider the following compound square of nine squares

-51	39	26	0	9	13	6	17	15	-6	0	4	21	35	12	-60	21	7	
54	-	-2	-5	4	-5	20	5	2	0	9	0	-34	19	28	2	13	8	
-5	1	2	3	17	18	-24	6	7	8	19	20	-42	0	15	3	20	40	
12	3	-1	63	-	-	18	12	8	6	-5	-3	72	-9	5	-9	-	0	
17	8	17	-42	22	14	12	3	12	-8	7	10	10	-1	4	46	-7	-16	
9	-5	-6	17	11	10	4	-7	-8	36	6	5	9	-8	-	54	12	-3	
2	53	45	-	33	34	59	31	34	-	137	24	25	74	49	31	-	45	28
-10	0	10	11	12	13	-44	15	14	16	17	18	-98	29	40	18	21	26	
-89	21	22	23	29	30	-	26	27	28	31	32	-	20	35	23	32	52	
143	-	10	-	-	-	149	-	-	-47	-	-	203	-	-	-62	-	-19	
21	22	41	33			12	13		19	22		33	16		37			
1	0	-1	22	12	2	-4	-5	-6	56	-3	-2	-6	-9	-	110	-	-28	
-11	-	101	-9	-	10	-16	-	-	120	-	-	-11	-	-	138	-8	-23	
48	7	79	-	10	14	105	-	68	-	1	5	120	3	65	-	22	8	
-44	1	25	16	27	11	-78	16	29	21	32	16	-	30	55	23	36	24	
-	33	6	9	4	85	-	38	11	14	6	87	-	32	19	9	7	107	
101						120	11	7	-93	27	-	194	-	4	-	9	-53	
134	2	-2	-36	5	-	140	-	-	-	-	-	-11	-	10	108	-		
-4	-	15	56	11	-	67	-9	-	-4	90	-4	-17	24	12	144	-18	-43	
3	8	-	113	-	6	73	-2	6	-	132	-	1	3	5	-	150	-7	

We note that the proposition ensures us that the following vectors form a basis of a subspace of the nullspace of the square:

$$(-33, -141, -123, 33, 141, 123, 33, 141, 123, -33, -141, -123, 0, 0, 0, 0, 0, 0)',$$

$$(-38, -837, 0, 38, 837, 0, -85, 837, 0, 85, -837, 0, 123, 0, 0, -123, 0, 0)',$$

$$(100, 513, 0, -100, -513, 0, -100, -636, 0, 100, 636, 0, 0, 123, 0, 0, -123, 0)',$$

$$(141, 759, -123, -141, -759, 123, -141, 0, 0, 141, 0, 0, 0, 123, 0, 0, 123)'. \quad .$$

REFERENCES

- [1] Ahmed, M., Algebraic Combinatorics of Magic Squares, Ph. D. Thesis, Mathematics Dep., University of California, Davis, 2004.
- [2] Al-Amerie, M., Msc. Thesis, Al-Albayt University, supervised by S. Al-Ashhab, S., 2007.

- [3] Al- Ashhab, S., Theory of Magic Squares, Royal Scientific Society, 2000.
- [4] Al-Ashhab, S., Mueller W., Semi Pandiagonal Magic 4×4 Squares, Results in Mathematics 44, (2003), p.p. (25-28).
- [5] Andress, W., Basic Properties of Pandiagonal Magic Squares, The American Mathematical Monthly, Vol. (67) No. (2), 1960, p.p. 143-152.
- [6] Anton, H., Elementary Linear Algebra, 7th Edition, Jon Wiley, New York, 1994.
- [7] Bellew, S., Counting the Number of Compound and Nasik Magic Squares, Mathematics Today, August (1997), p.p. 111-118.
- [8] Duijvestijn, A., Federico P., Compound Perfect Squares, the American Mathematical Monthly, Vol.89, No. 1, 1982, p.p. (15-32).
- [9] Hendricks, J., The Determinant of a Pandiagonal Magic Square of Order 4 is Zero, Journal of Recreational Mathematics, Vol. (21), 1989, p. p. 179-181.
- [10] Rosser, B. , Walker, R., On The Transformation Group for Diabolic Magic Squares of Order Four, Bulletin of the American Mathematical Society, Vol. (44), 1938, p.p. 416-420.
- [11] Ward, J., Vector Spaces of Magic Squares, Mathematics Magazine, Vol. (53), No. (2), March (1980), p. p. 108-111.