

# Based on a family of bi-univalent functions introduced through the $q$ -analogue of Noor integral operator

A. Akgül, B. Alçın

Kocaeli University  
e-mail:akgul@kocaeli.edu.tr  
Kocaeli University  
e-mail:bulentalcin41@gmail.com

Received 18 November 2021; Accepted 15 February 2022

## Abstract

*Recently,  $q$ -analogue of Noor integral operator and other special operators became importance in the field of Geometric Function Theory. In this study, by connecting this operators we introduced an interesting class of bi-univalent functions and obtained coefficient estimates for this new class. Also we present some relevant corollaries.*

**Keywords:** *analytic function, univalent and bi-univalent functions, subordination, coefficient bounds, integral operator.*

## 1 Introduction

Let  $\mathcal{A}$  indicates the family of analytic functions having form

$$f(z) = z + \sum_{n=2}^{\infty} a_n z^n \quad (1)$$

in the open unit disk  $\mathfrak{D} = \{z : |z| < 1, z \in \mathbb{C}\}$  and let  $\mathcal{S} = \{f \in \mathcal{A} : f \text{ is univalent in } \mathfrak{D}\}$ .

According the Koebe one-quarter theorem [14], the image of  $\mathfrak{D}$  under every function  $f$  from  $\mathcal{S}$  contains a disk of radius  $\frac{1}{4}$ . That is, every such univalent function has an inverse  $f^{-1}$  satisfying

$$f^{-1}(f(z)) = z \quad (z \in \mathfrak{D})$$

and

$$f(f^{-1}(w)) = w \quad \left( |w| < r_0(f), r_0(f) \geq \frac{1}{4} \right),$$

where

$$f^{-1}(w) = w - a_2 w^2 + (2a_2^2 - a_3) w^3 - (5a_2^3 - 5a_2 a_3 + a_4) w^4 + \dots \quad (2)$$

If  $f$  and  $f^{-1}$  are univalent, then we say that  $f$  is bi-univalent function in  $\mathfrak{D}$ . The class of bi-univalent functions defined in  $\mathfrak{D}$  is symbolized by  $\Sigma$ .

One can see important examples in the class in [27]. Although the functions  $\frac{z}{1-z}$ ,  $-\log(1-z)$ ,  $\frac{1}{2} \log\left(\frac{1+z}{1-z}\right)$  are in  $\Sigma$ , well known Koebe function is not in  $\Sigma$ . For example,  $z - \frac{z^2}{2}$  and  $\frac{z}{1-z^2}$  are in  $\mathcal{S}$  but not in  $\Sigma$  [27].

Given  $f, g \in \mathcal{A}$ ,  $f$  is said to be subordinate to  $g$ , symbolized

$$f(z) \prec g(z), \quad (3)$$

such that there is an analytic function  $w$  defined on  $\mathfrak{D}$  with

$$w(0) = 0 \quad \text{and} \quad |w(z)| < 1$$

fulfilling the following condition:

$$f(z) = g(w(z)).$$

The aforementioned subclasses of  $\Sigma$  were constructed and non-sharp estimates on the first two coefficients  $|a_2|$  and  $|a_3|$  in the Taylor-Maclaurin series expansion (1) were found in several recent studies (see [11, 12], [15], [27], [28, 29]). In very nearly, they have been followed by many works (see also [1], [8], [10, 13, 15, 16, 17], [19], [23], [22]).

Now, we give some basic definitions.

**Definition 1.1.** For  $q \in (0, 1)$ , the  $q$ -derivative of function  $f \in \mathcal{A}$  is defined by (see [18])

$$\partial_q f(z) = \frac{f(qz) - f(z)}{(q-1)z}, \quad z \neq 0 \quad (4)$$

and

$$\partial_q f(0) = f'(0).$$

Thus we have

$$\partial_q f(z) = 1 + \sum_{k=2}^{\infty} [k, q] a_k z^{k-1} \quad (5)$$

where  $[k, q]$  is given by

$$[k, q] = \frac{1 - q^k}{1 - q}, \quad [0, q] = 0 \quad (6)$$

and the  $q$ -fractional is defined by

$$[k, q]! = \begin{cases} \prod_{n=1}^k [n, q], & k \in \mathbb{N} \\ 1, & k = 0 \end{cases}.$$

Also, the  $q$ -generalized Pochhammer symbol for  $\mathfrak{p} \geq 0$  is given by

$$[\mathfrak{p}, q]_k = \begin{cases} \prod_{n=1}^k [\mathfrak{p} + n - 1, q], & k \in \mathbb{N} \\ 1, & k = 0 \end{cases}.$$

As  $q \rightarrow 1$ , then we get  $[k, q] \rightarrow k$ . Thus, by choosing the function  $g(z) = z^k$ , while  $q \rightarrow 1$ , then we obtain

$$\partial_q g(z) = \partial_q z^k = [k, q] z^{k-1} = g'(z),$$

where  $g'$  is the ordinary derivative.

Recently, function  $F_{q, \mu+1}^{-1}(z)$  is defined by Arif et al. [9] by

$$F_{q, \mu+1}^{-1}(z) * F_{q, \mu+1}(z) = z \partial_q f(z), \quad (\mu > -1)$$

where

$$F_{q, \mu+1}(z) = z + \sum_{k=2}^{\infty} \frac{[\mu + 1, q]_{k-1}}{[k - 1, q]!} z^k, \quad z \in \mathcal{D}. \quad (7)$$

To the series defined in (7) is convergent absolutely in  $\mathfrak{D}$ , by using the definition of  $q$ -derivative through convolution, let us explain the integral operator  $\zeta_q^\mu : \mathfrak{D} \rightarrow \mathfrak{D}$  by

$$\zeta_q^\mu f(z) = F_{q, \mu+1}^{-1}(z) * f(z) = z + \sum_{k=2}^{\infty} \phi_{k-1} a_k z^k, \quad (z \in \mathfrak{D}) \quad (8)$$

where

$$\phi_{k-1} = \frac{[k, q]!}{[\mu + 1, q]_{k-1}}. \quad (9)$$

From (8), one can readily have the identity

$$[\mu + 1, q] \zeta_q^\mu f(z) = [\mu, q] \zeta_q^{\mu+1} f(z) + q^\mu z \partial_q (\zeta_q^{\mu+1} f(z)).$$

We can state that

$$\zeta_q^0 f(z) = z \partial_q f(z), \quad \zeta_q' f(z) = f(z) \quad (10)$$

also

$$\lim_{q \rightarrow 1^-} \zeta_q^\mu f(z) = z + \sum_{k=2}^{\infty} \frac{k!}{(\mu+1)_{k-1}} a_k z^k.$$

This means that, by taking  $q \rightarrow 1^-$ , the operator defined in (8) reduces to the famous Noor integral operator given in ([20]). Moreover, for more detailed knowledge on the  $q$ -analogue of differential and integral operators, see the work of [20], [21]. Also for more details on the  $q$ -analogue of differential and integral operators, see the work of Aldweby and Darus (see[7]) (also see [1, 2, 3, 4, 8], [22]).

**Definition 1.2.** Let the functions  $h, p : \mathfrak{D} \rightarrow \mathbb{C}$  be so constrained that

$$\min \{ \Re(h(z)), \Re(p(z)) \} > 0$$

and

$$h(0) = p(0) = 1.$$

In this study, motivated by the work Altinkaya and Yalçın [6] (see also [5], [25], by the help of the aforementioned works we introduce a general new subclass  $\mathfrak{S}_{\Sigma_\gamma}^{\mu, q}$  of the function class  $\Sigma$  and obtain estimates of the coefficients  $|a_2|$  and  $|a_3|$  for functions in our new class. We state that through this paper,  $f, g$  are given by (1) and (2) respectively and  $\zeta_q^\mu$  is  $q$ -analogue of Noor integral operator.

## 2 The class $\mathfrak{S}_{\Sigma_\gamma}^{\mu, q}$

**Definition 2.1.** For  $f \in \Sigma, f \in \mathfrak{S}_{\Sigma_\gamma}^{\mu, q}$  if the following conditions are satisfied:

$$1 + \frac{1}{\tau} \left[ (1 - \gamma) \frac{\zeta_q^\mu f(z)}{z} + \gamma \partial_q (\zeta_q^\mu f(z)) - 1 \right] \in h(\mathfrak{D}), \quad (z \in \mathfrak{D}), \quad (11)$$

$$1 + \frac{1}{\tau} \left[ (1 - \gamma) \frac{\zeta_q^\mu g(w)}{w} + \gamma \partial_q (\zeta_q^\mu g(w)) - 1 \right] \in p(\mathfrak{D}), \quad (w \in \mathfrak{D}) \quad (12)$$

where  $\xi \geq 1, \tau \neq 0$ .

**Remark 2.2.** There are many choices of  $h, p$  and  $\mu, q, \gamma$  which would provide interesting subclasses of class  $\mathfrak{S}_{\Sigma_\gamma}^{\mu, q}$ . For example

1. For

$$h(z) = p(z) = \left( \frac{1+z}{1-z} \right)^\lambda, (0 < \lambda \leq 1)$$

it can be directly verified that the functions  $h(z)$  and  $p(z)$  satisfy the hypotheses of Definition 2. It is simply to see that  $f \in \Sigma$  is in  $\mathfrak{S}_{\Sigma_\gamma}^{\mu,q}(\lambda)$

if the following inequalities hold:

$$\left| \arg \left( 1 + \frac{1}{\tau} \left[ (1-\gamma) \frac{\zeta_q^\mu f(z)}{z} + \gamma \partial_q (\zeta_q^\mu f(z)) - 1 \right] \right) \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, z \in \mathfrak{D})$$

and

$$\left| \arg \left( 1 + \frac{1}{\tau} \left[ (1-\gamma) \frac{\zeta_q^\mu g(w)}{w} + \gamma \partial_q (\zeta_q^\mu g(w)) - 1 \right] \right) \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, w \in \mathfrak{D}).$$

In the case of  $h(z) = p(z) = \left( \frac{1+z}{1-z} \right)^\lambda$  where  $(0 < \lambda \leq 1)$ , the class  $\mathfrak{S}_{\Sigma_\gamma}^{\mu,q}$  reduces following new also well known subclasses:

i) Upon setting  $q \rightarrow 1^-$  and it can be directly verified that the functions  $h(z)$  and  $p(z)$  satisfy the hypotheses of Definition 2. it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_\gamma}^{\mu,1}(\lambda) = \mathfrak{S}_{\Sigma_\gamma}^\mu(\lambda)$$

if the following inequalities hold:

$$\left| \arg \left( 1 + \frac{1}{\tau} \left[ (1-\gamma) \frac{\zeta^\mu f(z)}{z} + \gamma (\zeta^\mu f(z))' - 1 \right] \right) \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, z \in \mathfrak{D})$$

and

$$\left| \arg \left( 1 + \frac{1}{\tau} \left[ (1-\gamma) \frac{\zeta^\mu g(w)}{w} + \gamma (\zeta^\mu g(w))' - 1 \right] \right) \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, w \in \mathfrak{D}).$$

ii) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{\mu,1}(\lambda) = \mathfrak{S}_{\Sigma}^\mu(\lambda)$$

if the following inequalities hold:

$$\left| \arg \left[ (1-\gamma) \frac{\zeta^\mu f(z)}{z} + \gamma (\zeta^\mu f(z))' \right] \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, z \in \mathfrak{D})$$

and

$$\left| \arg \left[ (1-\gamma) \frac{\zeta^\mu g(z)}{z} + \gamma (\zeta^\mu g(z))' \right] \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, z \in \mathfrak{D}).$$

- iii) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$  and  $\mu = 1$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{1,1}(\lambda) = \mathfrak{S}_{\Sigma_1}(\lambda)$$

if the following inequalities hold:

$$\left| \arg \left[ (1 - \gamma) \frac{f(z)}{z} + \gamma (f(z))' \right] \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D})$$

and

$$\left| \arg \left[ (1 - \gamma) \frac{g(w)}{w} + \gamma (g(w))' \right] \right| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D})$$

- iv) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$ ,  $\gamma = 1$  and  $\mu = 1$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{1,1}(\lambda) = \mathfrak{S}_{\Sigma}(\lambda) = \mathcal{H}_{\Sigma}^{\lambda}$$

The class  $\mathcal{H}_{\Sigma}^{\lambda}$  was introduced by Srivastava et al. [27] The function  $f \in \mathcal{H}_{\Sigma}^{\lambda}$ , if the following inequalities hold:

$$|\arg f'(z)| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D})$$

and

$$|\arg g'(z)| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D}).$$

- v) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$ ,  $\gamma = 1$  and  $\mu = 0$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{\circ,1}(\lambda) = \mathfrak{S}_{\Sigma_1}^{\circ}(\lambda)$$

if the following inequalities hold:

$$|\arg (z\partial f(z))'| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D})$$

and

$$|\arg (w\partial g(w))'| < \frac{\lambda\pi}{2} \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D}).$$

2. For

$$h(z) = p(z) = \frac{1 + (1 - 2\beta)z}{1 - z}, \quad (0 \leq \beta < 1)$$

it can be directly verified that the functions  $h(z)$  and  $p(z)$  satisfy the hypotheses of Definition 2. It is simply to see that  $f \in \Sigma$  is in  $\mathfrak{S}_{\Sigma_{\gamma}}^{\mu,q}(\beta)$ , if the following inequalities hold:

$$\Re \left( 1 + \frac{1}{\tau} \left[ (1 - \gamma) \frac{\zeta_q^{\mu} f(z)}{z} + \gamma \partial_q (\zeta_q^{\mu} f(z)) - 1 \right] \right) > \beta \quad (0 \leq \beta < 1, \quad 0 < \alpha \leq 1, \quad z \in \mathfrak{D})$$

and

$$\Re \left( 1 + \frac{1}{\tau} \left[ (1 - \gamma) \frac{\zeta_q^\mu g(w)}{w} + \gamma \partial_q (\zeta_q^\mu g(w)) - 1 \right] \right) > \beta \quad (0 \leq \beta < 1, 0 < \alpha \leq 1, w \in \mathfrak{D}) .$$

In the case of  $h(z) = p(z) = \frac{1+(1-2\beta)z}{1-z}$  where  $(0 \leq \beta < 1)$ , the class  $\mathfrak{S}_{\Sigma_\gamma}^{\mu,q}$  reduces following new also well known subclasses:

- i) Upon setting  $q \rightarrow 1^-$  and it can be directly verified that the functions  $h(z)$  and  $p(z)$  satisfy the hypotheses of Definition 2. it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_\gamma}^{\mu,1}(\beta) = \mathfrak{S}_{\Sigma_\gamma}^\mu(\beta)$$

if the following inequalities hold:

$$\Re \left( 1 + \frac{1}{\tau} \left[ (1 - \gamma) \frac{\zeta^\mu f(z)}{z} + \gamma (\zeta^\mu f(z))' - 1 \right] \right) > \beta \quad (0 \leq \beta < 1, z \in \mathfrak{D})$$

and

$$\Re \left( 1 + \frac{1}{\tau} \left[ (1 - \gamma) \frac{\zeta_q^\mu g(w)}{w} + \gamma \partial_q (\zeta_q^\mu g(w)) - 1 \right] \right) > \beta \quad (0 \leq \beta < 1, w \in \mathfrak{D}) .$$

- ii) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{\mu,1}(\beta) = \mathfrak{S}_{\Sigma}^\mu(\beta)$$

if the following inequalities hold:

$$\Re \left( (1 - \gamma) \frac{\zeta^\mu f(z)}{z} + \gamma (\zeta^\mu f(z))' \right) > \beta \quad (0 \leq \beta < 1, z \in \mathfrak{D})$$

$$\Re \left( (1 - \gamma) \frac{\zeta^\mu g(z)}{z} + \gamma (\zeta^\mu g(z))' \right) > \beta \quad (0 \leq \beta < 1, w \in \mathfrak{D}) .$$

- iii) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$  and  $\mu = 1$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{1,1}(\beta) = \mathfrak{S}_{\Sigma_1}(\beta)$$

if the following inequalities hold:

$$\Re \left( (1 - \gamma) \frac{f(z)}{z} + \gamma (f(z))' \right) > \beta \quad (0 \leq \beta < 1, z \in \mathfrak{D})$$

and

$$\Re \left( (1 - \gamma) \frac{g(w)}{w} + \gamma (g(w))' \right) > \beta \quad (0 \leq \beta < 1, w \in \mathfrak{D})$$

- iv) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$ ,  $\gamma = 1$  and  $\mu = 1$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{1,1}(\beta) = \mathfrak{S}_{\Sigma}(\beta) = \mathcal{H}_{\Sigma}^{\beta}$$

The class  $\mathcal{H}_{\Sigma}^{\beta}$  was introduced by Srivastava et al. [27] The function  $f \in \mathcal{H}_{\Sigma}^{\beta}$ , if the following inequalities hold:

$$\Re(f'(z)) > \beta \quad (0 < \beta \leq 1, \quad z \in \mathfrak{D})$$

and

$$\Re(g'(z)) > \beta \quad (0 < \beta \leq 1, \quad z \in \mathfrak{D}).$$

- v) Upon setting  $q \rightarrow 1^-$ , for  $\tau = 1$ ,  $\gamma = 1$  and  $\mu = 0$ , it is simply to see that  $f \in \Sigma$  is in

$$\mathfrak{S}_{\Sigma_1}^{\circ,1}(\lambda) = \mathfrak{S}_{\Sigma_1}^{\circ}(\lambda)$$

if the following inequalities hold:

$$|\arg(z\partial f(z))'| > \beta \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D})$$

and

$$|\arg(w\partial g(w))'| > \beta \quad (0 < \lambda \leq 1, \quad z \in \mathfrak{D}).$$

We need the following lemma to derive our main result.

**Theorem 2.3.** *Let  $f$  given by (1) be in the class  $\mathfrak{S}_{\Sigma_{\gamma}}^{\mu,q}$ . Then*

$$|a_2| \leq \min \left\{ \sqrt{\frac{(|h'(0)|^2 + |p'(0)|^2)\tau^2}{(1+\gamma q)^2 \phi_1^2}}, \sqrt{\frac{(|h''(0)| + |p''(0)|)\tau}{4(1+\gamma q + \gamma q^2)\phi_2}} \right\} \quad (13)$$

and

$$|a_3| \leq \min \left\{ \begin{array}{l} \frac{(|h'(0)|^2 + |p'(0)|^2)\tau^2}{(1+\gamma q)^2 \phi_1^2} + \frac{|h''(0)| + |p''(0)|}{2(1+\gamma q + \gamma q^2)\phi_2} |\tau|, \\ \frac{|h''(0)|}{(1+\gamma q + \gamma q^2)\phi_2} |\tau| \end{array} \right. \quad (14)$$

*Proof.* Let  $f \in \mathfrak{S}_{\Sigma_{\gamma}}^{\mu,q}$ . Firstly, we write the argument inequalities in (11) and (12) and in their equivalent forms as follows:

$$\begin{aligned} 1 + \frac{1}{\tau} \left[ (1-\gamma) \frac{\zeta_q^{\mu} f(z)}{z} + \gamma \partial_q(\zeta_q^{\mu} f(z)) - 1 \right] &= h(z) \\ 1 + \frac{1}{\tau} \left[ (1-\gamma) \frac{\zeta_q^{\mu} g(w)}{w} + \gamma \partial_q(\zeta_q^{\mu} g(w)) - 1 \right] &= p(w) \end{aligned} \quad (15)$$

where  $h(z)$  and  $p(w)$  satisfy the conditions of Definiton 2. Furthermore, the functions  $h(z)$  and  $p(w)$  have the following Taylor-Maclaurin series expansions:

$$\begin{aligned} h(z) &= 1 + h_1z + h_2z^2 + \dots, \\ p(w) &= 1 + p_1w + p_2w^2 + \dots, \end{aligned} \quad (16)$$

respectively. Thus, upon comparing the corresponding coefficients in (15), we get

$$\frac{1}{\tau} (1 + \gamma q) \phi_1 a_2 = h_1 \dots \quad (17)$$

$$\frac{1}{\tau} (1 + \gamma q + \gamma q^2) \phi_2 a_3 = h_2 \dots \quad (18)$$

and

$$-\frac{1}{\tau} (1 + \gamma q) \phi_1 a_2 = p_1.. \quad (19)$$

$$\frac{1}{\tau} [(1 + \gamma q + \gamma q^2) \phi_2 (2a_2^2 - a_3)] = p_2 \dots \quad (20)$$

From (17) and (19), we have

$$h_1 = -p_1 \dots \quad (21)$$

and

$$2 \left[ \frac{1}{\tau} (1 + \gamma q) \phi_1 \right]^2 a_2^2 = h_1^2 + p_1^2. \dots \quad (22)$$

Also from (18) and (20), we get

$$\frac{2}{\tau} [(1 + \gamma q + \gamma q^2) \phi_2] a_2^2 = h_2 + p_2 \dots \quad (23)$$

which gives us the desired estimate on  $|a_2|$  as asserted in (13).

Next, in order to find the bound on  $|a_3|$ , by subtracting (20) from (18), we obtain

$$a_3 - a_2^2 = \frac{h_2 - p_2}{2(1 + \gamma q + \gamma q^2) \phi_2} \tau \quad (24)$$

Therefore, in view of (22) and (23), we have

$$a_3 = \frac{h_2 - p_2}{2(1 + \gamma q + \gamma q^2) \phi_2} \tau + \frac{(h_1^2 + p_1^2)}{2(1 + \gamma q)^2 \phi_1^2} \tau^2$$

which completes the proof of Theorem.  $\square$

### 3 Corollaries and Consequences

**Corollary 3.1.** *If we let*

$$h(z) = p(z) = \left( \frac{1+z}{1-z} \right)^\lambda = 1 + 2\lambda z + 2\lambda^2 z^2 + \dots \quad (0 < \lambda \leq 1),$$

then inequalities (13) and (14) become

$$|a_2| \leq \min \left\{ \frac{2\lambda|\tau|}{(1+\gamma q)\phi_1}, \sqrt{\frac{2\lambda\tau}{(1+\gamma q+\gamma q^2)\phi_2}} \right\} = \sqrt{\frac{2\lambda\tau}{(1+\gamma q+\gamma q^2)\phi_2}} \quad (25)$$

and

$$|a_3| \leq \min \left\{ \frac{4\lambda^2\tau^2}{(1+\gamma q)^2\phi_1^2} + \frac{2\lambda|\tau|}{(1+\gamma q+\gamma q^2)\phi_2}, \frac{2\lambda|\tau|}{(1+\gamma q+\gamma q^2)\phi_2} \right\}. \quad (26)$$

**Corollary 3.2.** *If we let*

$$h(z) = p(z) = \frac{1+(1-2\beta)z}{1-z} = 1+2(1-\beta)z+2(1-\beta)z^2+\dots \quad (0 \leq \beta < 1),$$

then inequalities (13) and (14) become

$$|a_2| \leq \min \left\{ \frac{2(1-\beta)|\tau|}{(1+\gamma q)\phi_1}, \sqrt{\frac{2(1-\beta)\tau}{(1+\gamma q+\gamma q^2)\phi_2}} \right\}.$$

and

$$|a_3| \leq \min \left\{ \frac{4(1-\beta)^2\tau^2}{(1+\gamma q)^2\phi_1^2} + \frac{2(1-\beta)|\tau|}{(1+\gamma q+\gamma q^2)\phi_2}, \frac{2(1-\beta)|\tau|}{(1+\gamma q+\gamma q^2)\phi_2} \right\}.$$

An urgent and important corollary of the Theorem for  $q \rightarrow 1^-$  is asserted by following Corollary.

**Corollary 3.3.** *If  $f \in \mathfrak{S}_{\Sigma_\gamma}^{\mu,q}$ , then*

$$|a_2| \leq \min \left\{ \sqrt{\frac{(|h'(0)|^2+|p'(0)|^2)\tau^2}{(1+\gamma)^2\phi_1^2}}, \sqrt{\frac{(|h''(0)|+|p''(0)|)\tau}{4(1+2\gamma)\phi_2}} \right\} \quad (27)$$

and

$$|a_3| \leq \min \left\{ \begin{array}{l} \frac{(|h'(0)|^2+|p'(0)|^2)}{(1+\gamma)^2\phi_1^2}\tau^2 + \frac{|h''(0)|+|p''(0)|}{2(1+2\gamma)\phi_2}|\tau|, \\ \frac{|h''(0)|}{(1+2\gamma)\phi_2}|\tau| \end{array} \right\}. \quad (28)$$

**Corollary 3.4.** *If we let*

$$h(z) = p(z) = \left( \frac{1+z}{1-z} \right)^\lambda = 1 + 2\lambda z + 2\lambda^2 z^2 + \dots \quad (0 < \lambda \leq 1),$$

*then inequalities (27) and (28) become*

$$|a_2| \leq \min \left\{ \frac{2\lambda|\tau|}{(1+\gamma)\phi_1}, \sqrt{\frac{2\lambda\tau}{(1+2\gamma)\phi_2}} \right\} = \sqrt{\frac{2\lambda\tau}{(1+2\gamma)\phi_2}} \quad (29)$$

*and*

$$|a_3| \leq \min \left\{ \frac{4\lambda^2\tau^2}{(1+\gamma)^2\phi_1^2} + \frac{2\lambda|\tau|}{(1+2\gamma)\phi_2}, \frac{2\lambda|\tau|}{(1+2\gamma)\phi_2} \right\}. \quad (30)$$

*By choosing  $\gamma = 1$  in inequalities (29) and (30) we get*

$$|a_2| \leq \min \left\{ \frac{\lambda|\tau|}{\phi_1}, \sqrt{\frac{2\lambda\tau}{3\phi_2}} \right\} = \sqrt{\frac{2\lambda\tau}{3\phi_2}}$$

*and*

$$|a_3| \leq \min \left\{ \frac{\lambda^2\tau^2}{\phi_1^2} + \frac{2\lambda|\tau|}{3\phi_2}, \frac{2\lambda|\tau|}{3\phi_2} \right\}.$$

**Corollary 3.5.** *If we let*

$$h(z) = p(z) = \frac{1 + (1-2\beta)z}{1-z} = 1 + 2(1-\beta)z + 2(1-\beta)z^2 + \dots \quad (0 \leq \beta < 1),$$

*then inequalities (27) and (28) become*

$$|a_2| \leq \min \left\{ \frac{2(1-\beta)|\tau|}{(1+\gamma)\phi_1}, \sqrt{\frac{2(1-\beta)\tau}{(1+2\gamma)\phi_2}} \right\}. \quad (31)$$

*and*

$$|a_3| \leq \min \left\{ \frac{4(1-\beta)^2\tau^2}{(1+\gamma)^2\phi_1^2} + \frac{2(1-\beta)|\tau|}{(1+2\gamma)\phi_2}, \frac{2(1-\beta)|\tau|}{(1+2\gamma)\phi_2} \right\}. \quad (32)$$

*By choosing  $\gamma = 1$  in inequalities (31) and (32) we get*

$$|a_2| \leq \min \left\{ \frac{(1-\beta)|\tau|}{\phi_1}, \sqrt{\frac{2(1-\beta)\tau}{3\phi_2}} \right\}. \quad (33)$$

*and*

$$|a_3| \leq \min \left\{ \frac{(1-\beta)^2\tau^2}{\phi_1^2} + \frac{2(1-\beta)|\tau|}{3\phi_2}, \frac{2(1-\beta)|\tau|}{3\phi_2} \right\}. \quad (34)$$

## 4 Conclusion

In this paper, we defined a new subclasses of bi-univalent functions in the open unit disk  $D$ , by using  $q$ -analogue of Noor integral operator. For the functions belonging to this subclass, the estimates of second and third Taylor–Maclaurin coefficients are obtained. Also, some interesting corollaries and remarks are discussed and relevant connections are made.

## 5 Open Problem

For the subclass  $\mathfrak{S}_{\Sigma_{\gamma}}^{\mu, q}$ , it is interesting to obtain the upper bounds of the Fekete–Szegő inequality, and the bounds of the Hankel determinants. Also, to examine the estimates of second and third Taylor–Maclaurin coefficients  $m$ -fold symmetric bi-univalent functions.

## References

- [1] Akgül, A. and Sakar, F. M., "A certain subclass of bi-univalent analytic functions introduced by means of the  $q$ -analogue of Noor integral operator and Horadam polynomials", *Turkish J. Math.*, 43(5) 2019 2275-2286.
- [2] Akgül, A., "A new approach to the bi-univalent analytic functions connected with  $q$ -analogue of Noor integral operator", *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.*, 70(2) (2021) 940-949.
- [3] Akgül, A., "On a special second order differential subordination using  $q$ -analogue of Noor integral operator", *Tbil. Math. J.* Special Issue (DECOP - 2020) (2022) 109-123.
- [4] Akgül, A. and Altınkaya, Ş., "Coefficient estimates associated with a new subclass of bi-univalent functions", *Acta Univ. Apul.*, 52(4) (2017) 121-128.
- [5] Ş. Altınkaya, S. Yalçın, Coefficient Estimates for Two New Subclasses of Bi-univalent Functions, *Acta Univ. Apul*, 43 (2015) 53-63.
- [6] Ş. Altınkaya, S. Yalçın, "Coefficient Estimates for Two New Subclasses of Bi-univalent Functions with respect to Symmetric Points", *Journal of Function Spaces*, Article ID 145242, (2015) 5 pp.
- [7] Aldweby, H. and Darus, M. "A subclass of harmonic univalent functions associated with  $q$ -analogue of Dziok–Srivastava operator", *ISRN Mathematical Analysis*, vol. 2013, (2013) Article ID 382312, 6 pages.

- [8] Altinkaya, Ş., "Inclusion properties of Lucas polynomials for bi-univalent functions introduced through the  $q$ -analogue of Noor integral operator. *Turkish J. Math.* 43 (2) (2019) 620-629.
- [9] Arif, M., Ul Haq M, Liu J-L. "Subfamily of univalent functions associated with  $q$ -analogue of Noor integral operator", *Journal of Function Spaces* (2018) (2018) 1-5.
- [10] Bulut, S., "Coefficient estimates for a class of analytic and bi-univalent functions", *Novi Sad J. Math.* 43 (2013)(2) 59-65.
- [11] Brannan, D. A. and Taha, T. S., "On some classes of bi-univalent functions", in: *S.M. Mazhar, A. Hamoui, N.S. Faour (Eds.), Math. Anal. and Appl., Kuwait; February 18-21, 1985, in: KFAS Proceedings Series, vol. 3, Pergamon Press, Elsevier Science Limited, Oxford, 1988, pp. 53-60. see also Studia Univ. Babeş-Bolyai Math.* 31 (2) (1986) 70-77.
- [12] Brannan, D. A. and Taha, T. S., "On some classes of bi-univalent functions", *Studia Universitatis Babes -Bolyai Mathematica*, 31(2) (1986) 70-77.
- [13] Çağlarlar M. , Orhan, H. and Yağmur, N., "Coefficient bounds for new subclasses of bi-univalent functions", *Filomat*, 27(7) (2013) 1165-1171.
- [14] Duren, P. L., *Univalent Functions*, vol. 259 of Grundlehren der Mathematischen Wissenschaften, Springer, New York, NY, USA, 1983.
- [15] Frasin, B. A. and Aouf, M. K., "New subclasses of bi-univalent functions", *Appl. Math. Lett.*, 24(9) (2011) 1569-1573.
- [16] Goyal, S. P. and Goswami, P., "Estimate for initial Maclaurin coefficients of bi-univalent functions for a class defined by fractional derivatives", *J. Egyptian Math. Soc.*, 20(3) ( 2012) 179-182.
- [17] T. Hayami and S. Owa, "Coefficient bounds for bi-univalent functions", *Pan-American Mathematical Journal*, 22(4) (2012) 15-26.
- [18] Jackson, F.H., "On  $q$ -functions and a certain difference operator" *Trans. R. Soc. Edinb.* 46 (1908) 253-281.
- [19] Orhan, H., Magesh, N., Balaji, V. K., "Initial coefficient bounds for a general class of bi-univalent functions" *Filomat*, 29(6) (2015) 1259-1267.
- [20] Noor K. I., "On new classes of integral operators", *J. Natur. Geom.* 16 (1999) 71-80.

- [21] Noor KI, Noor MA., "On integral operators" *J. Math. Anal. Appl.*, 238(2) (1999 Oct 15) 341-52.
- [22] Noor, K. I., Altinkaya, Ş., Yalçın, S., "Coefficient inequalities of analytic functions equipped with conic domains involving  $q$ -analogue of Noor integral operator", *Tbil. Math. J.*, 14(1) (2021) 1-14.
- [23] Porwal S. and Darus M., "On a new subclass of bi-univalent functions", *J. Egyptian Math. Soc.*, 21(3) (2013) 190–193.
- [24] Rogosinski, W., "On the coefficients of subordinate functions", *Proc. Lond. Math. Soc.*, 48(2) (1943) 48-82.
- [25] Sakar, F. M. and Akgül, A., "Based on a family of bi-univalent functions introduced through the Faber polynomial expansions and Noor integral operator", *AIMS Mathematics*, 7(4) (2022) 5146-5155.
- [26] Sălăgean G. S., "Subclasses of univalent functions", in *Complex Analysis-Fifth Romanian-Finnish Seminar, Part 1* (Bucharest, 1981), vol. 1013 of Lecture Notes in Mathematics, pp. 362-372, Springer, Berlin, Germany, 1983.
- [27] Srivastava, H.M., Mishra A. K., and Gochhayat P., "Certain subclasses of analytic and bi-univalent functions", *Appl. Math. Lett.*, 23(10) (2010) 1188-1192.
- [28] Xu, Q. H., Gui, Y. C., and Srivastava, H. M., "Coefficient estimates for a certain subclass of analytic and bi-univalent function", *Appl. Math. Lett.*, 25(6)(2012) 990-994.
- [29] Xu, Q. H., Xiao, H. G. and Srivastava, H. M., "A certain general subclass of analytic and bi-univalent functions and associated coefficient estimate problems", *Appl. Math. Comput.*, 218(23) (2012) 11461-11465.